

Forest Management



A Resource Management Strategy of the California Water Plan

California Department of Water Resources

July 29, 2016



Table of Contents

Forest Management	1
Forest Ownership and Management in California	1
Effects of Forest Management on Water Supply	4
Vegetation Management for Water Supply	5
Fuels/Fire Management	6
Wildfire Impacts on Watershed Resources	6
Fuel Treatments to Reduce Wildfire Impacts on Watershed Resources	8
Management Strategies to Reduce Adverse Impacts Associated with Wildfire	9
Recommendations	10
Road Management	10
Riparian Forests	11
Meadow Restoration and Groundwater	13
Urban Forestry	15
Urban Watershed Forestry	15
Tree Cover and Watershed Benefits	15
Stormwater Runoff	15
Structural Soils	17
Quantifying Benefits	17
Recommendations	17
Forest Loss and Fragmentation	18
Recommendation	18
Illegal Marijuana Cultivation	18
Climate Change	20
Adaptation	20
Mitigation	20
Potential Costs	21
Vegetation Management	21
Fuels/Fire Management	21
Road Management	21
Riparian Forests	21
Meadow Groundwater Storage	21
Urban Forestry	22
Illegal Marijuana Cultivation	22
Major Implementation Issues	22
Information Needs	22
Coordination Needs	23
Limited Funding for Forest Watershed Restoration	23
Regulatory Requirements	24
Overarching Recommendations	25
Monitoring and Research	25
Coordination	25
Regulatory Requirements	26
Beneficial Forest Management in Areas with Commercial-Scale Marijuana Cultivation	26
Special Author’s Note	27
References	27
References Cited	27
Additional References	42

Tables

Table 1 Acres of Forest Land by Ownership in California	2
Table 2 Watershed Benefits of Urban Forest Cover	16

Acronyms and Abbreviations

AB 32	climate change legislation
af	acre-foot, acre-feet
ARB	California Air Resources Board
BAER	Burned Area Emergency Response
BLM	U.S. Bureau of Land Management
BMP	best management practice
BOF	California State Board of Forestry and Fire Protection
Cal EMA	California Emergency Management Agency
Cal EPA	California Environmental Protection Agency
CAL FIRE	California Department of Forestry and Fire Protection
CEQA	California Environmental Quality Act
CO ₂	carbon dioxide
EPA	U.S. Environmental Protection Agency
FPR	Forest Practice Rule
GHG	greenhouse gas
HFI	Healthy Forest Initiative
LRMP	Land and Resource Management Plan
NEPA	National Environmental Policy Act
NFP	National Fire Plan
NPS	National Park Service
RWQCB	regional water quality control board
SWP	State Water Project

Forest Management

SWRCB	State Water Resources Control Board
THP	timber harvesting plan
TMDL	total maximum daily load
USDA Forest Service	U.S. Department of Agriculture Forest Service
USGS	U.S. Geological Survey
WDR	Waste Discharge Requirements or Waivers of Waste Discharge Requirements

Forest Management

Forest lands in California, the majority of which are in the middle to high elevation foothills and mountains, produce a diverse array of resources such as water, timber, native vegetation, fish, wildlife, and livestock, and outdoor recreation. However, the water produced by these forests has economic value that equals or exceeds that of any other forest resource (Krieger 2001; California Department of Forestry and Fire Protection 2003). Most of California's major rivers and a substantial portion of its runoff originate in these forests; therefore, most of California's major water development projects are tied strongly to forested watersheds.

Forest management activities can affect water quantity and quality. This strategy focuses on forest management activities, on both public and privately-owned forest lands, whose goals specifically include improvement of the availability and quality of water for downstream users.

Water rights for groundwater in most areas of California are assigned to overlying landowners and reasonable use is unregulated. In contrast, surface water rights, which are managed and enforced by the State Water Resources Control Board (SWRCB), are a complicated mixture of riparian, appropriative, and adjudicated rights. The U.S. Department of Agriculture Forest Service (USDA Forest Service) uses federal reserved, appropriative, riparian, and overlying adjudicated water rights to manage forest lands. A large percentage of water flowing from forests is appropriated by state and federal water projects, municipal water agencies, irrigation districts and hydropower companies, many of which are fully appropriated. A list of fully appropriated stream systems for California is available on the SWRCB web site: http://www.waterboards.ca.gov/waterrights/water_issues/programs/fully_appropriated_streams/.

Water quality in California is protected by the SWRCB and nine regional water quality control boards (RWQCBs). The RWQCBs regulate compliance with the federal Clean Water Act through designation of beneficial uses, development of numeric and narrative water quality objectives, water quality control policies, basin plans, basin plan prohibitions, issuance of various types of permits, and enforcement actions. The SWRCB prepares lists of impaired water bodies every two years, as required by Section 303(d) of the Clean Water Act. In addition to listings of threatened or endangered fish and wildlife species by the California Department of Fish and Wildlife, the U.S. Fish and Wildlife Service, and the National Marine Fisheries Service, these impaired water body listings have greatly influenced forestry practices on non-federal lands during the past decade (Cafferata et al. 2007a).

Forest Ownership and Management in California

California has more than 30 million acres of forested lands (California Department of Forestry and Fire Protection 2003; Christensen 2008), which are located primarily in the major mountains of the Coast Ranges, Klamath Mountains, Cascade Range, and the Sierra Nevada. Forest lands in California are owned and managed by a wide array of federal, state, tribal, and local agencies, non-governmental organizations, and private companies, families, and individuals (Table 1), each of whom has a different forest management strategy with different goals and constraints.

Table 1 Acres of Forest Land by Ownership in California

Landowner	Acres ^a	Percentage
Private non-corporate	8,448,000	22.5
Private corporate	4,719,000	12.6
County	330,000	0.9
State	726,000	1.9
USFS ^b	20,166,000	53.7
BLM	1,650,000	4.4
NPS	1,287,000	3.4
Other federal	231,000	0.6
Total	37,557,000	100.0

Source: California Department of Water Resources 2011, Christensen, et al. 2008.

Notes:

^a Acres reported are "real" forest rather than total ownership.

^b USDA Forest Service 2008.

The largest public forest landowner in the state is the USDA Forest Service, which owns and manages 18 National Forests in California. California's National Forests were established under the Organic Act of 1897, which specifically states that a primary purpose of these lands is to "secure favorable conditions of water flow." U.S. Secretary of Agriculture Tom Vilsach emphasized the role of the USDA Forest Service in protecting water sources in his remarks made on August 14, 2009:

We must work and must be committed to a shared vision, a vision that conserves our forests and the vital resources important to our survival while wisely respecting the need for a forest economy that creates jobs and vibrant rural communities. Our shared vision must begin with a complete commitment to restoration. Restoration, for me, means managing forest lands first and foremost to protect our water sources while making our forests far more resilient to climate change.

The USDA Forest Service Pacific Southwest Region manages roughly 20 million acres in California for multiple uses including, among other things, timber and livestock production, mineral production, and outdoor recreation (USDA Forest Service 2007). Despite their name, these National Forests include a wide variety of ecological communities, including subalpine and montane forests, alpine shrublands, chaparral, desert, and wetlands. Timber on National Forests is produced through commercial timber sales to private contractors and livestock are grazed under a permit system. Environmental issues related to resource management on National Forests are addressed under the National Environmental Policy Act (NEPA). Resource management on each National Forest is guided by a Land and Resource Management Plan (LRMP), which is revised and updated roughly every fifteen years. The content and format of LRMPs is governed by national planning rules, which are also revised periodically, with the most recent planning rule completed in 2011. All future LRMPs will emphasize sustainability, restoration, and forest health.

The U.S. Bureau of Land Management (BLM) manages 1,650,000 acres of forest in the state (USDA Forest Service 2008), primarily in the North Coast region and the Modoc Plateau. The BLM is a multiple-

use land management agency that produces timber through commercial sales and manages livestock grazing through a permit system. Environmental issues related to resource management on public lands administered by the BLM are addressed under NEPA.

The National Park Service (NPS) manages 1,287,000 acres of forest in 23 units in California (USDA Forest Service 2008). Unlike the USDA Forest Service and BLM, the NPS is not a multiple-use management agency, but instead has a mission to preserve natural and cultural resources specifically for public enjoyment and scientific purposes. Commercial timber harvests and livestock grazing are not allowed in national parks, although vegetation may be managed for forest health and fire protection purposes. Pack stock grazing is allowed when permitted.

Commercial timberlands (forests used or suitable for producing timber) comprise 16.6 million acres of forest land across the state (California Department of Forestry and Fire Protection 2003), nearly half of which is in non-federal ownership. More than five million acres are zoned for timber production and are primarily managed by large, industrial landowners (California Department of Forestry and Fire Protection 2003), with the remaining non-federal timberlands owned primarily by small non-industrial landowners with a wide range of management objectives. State Demonstration Forests include about 71,000 acres statewide (California Department of Forestry and Fire Protection 2013). Timber harvesting on non-federal forest lands is regulated by the California State Board of Forestry and Fire Protection (BOF) and the California Department of Forestry and Fire Protection (CAL FIRE). The BOF adopts regulations that CAL FIRE has enforced on the ground since 1975. Timber production is the primary use of privately-held forests, but some company-owned forest lands are used for livestock grazing and permitted outdoor recreation, including fishing and hunting. In addition, with the passage of recent climate change legislation (AB 32), some forests are likely to be managed to enhance carbon sequestration and provide offsets to greenhouse gas (GHG) emissions.

Urban forestry, although geographically distinct from wildland forests, offers important benefits for water resources and mitigation of climate change. Urban forests are managed by municipal parks and public works departments, as well as by many private organizations and individuals. Trees in urban environments provide more than just aesthetic benefits, including interception of rainfall, reduction of urban runoff, and energy-efficient shade during hot weather.

Forest management agencies also have responsibilities for protection of water quality and beneficial uses, such as aquatic habitat. The USDA Forest Service, CAL FIRE, and BOF have been designated by the SWRCB as water quality management agencies for lands that they administer or regulate, and have all implemented water quality management plans that have been certified by the SWRCB. These water quality management programs incorporate best management practices (BMPs) or Forest Practice Rules (FPRs) that are designed to prevent adverse impacts to water quality from forest management activities, and include monitoring programs to evaluate BMP/FPR implementation and effectiveness. The USDA Forest Service water quality program also includes restoration of “legacy” sources of pollution.

Extensive monitoring of California’s FPRs for protecting water quality on non-federal timberlands was conducted from 1996 through 2004 by two State programs — one using independent contractors acting as third party auditors to collect field data, and one using CAL FIRE forest practice inspectors (Cafferata and Munn 2002; Ice et al. 2004; Brandow et al. 2006). Together, these projects inspected more than 600 randomly selected timber harvesting plans (THPs) that had gone through one or more over-wintering

periods after the completion of logging. Both projects found that hillslope surface erosion features were almost always associated with improperly implemented forest practice rules on forest roads and at watercourse crossings, and that watercourse and lake protection zones (buffer strips) retained high levels of post-harvest canopy and surface cover, which prevented harvest-related erosion. Approximately 20 percent of stream crossings were found to have significant problems with both forest practice rule implementation and effectiveness. Overall, California forest practice rule implementation rates have been found to be among the highest of any of the Western states, and when properly implemented, these practices have been found to be highly effective in preventing hillslope erosion features (Ice et al. 2004; Council of Western State Foresters 2007; Ice et al. 2010).

The USDA Forest Service reported on monitoring data collected from 2003 through 2007 at roughly 2,900 randomly located sites to evaluate the implementation and effectiveness of its water quality BMPs on National Forests (USDA Forest Service 2009). The BMP Evaluation Program uses 29 different onsite monitoring protocols to evaluate BMP implementation and effectiveness, with the majority related to timber and engineering practices. Overall, 86 percent of the BMPs evaluated were rated as correctly and fully implemented, and 93 percent of these were rated as effective in protecting water quality, comparable to results documented on private timber lands. Among all evaluations, 98 percent were found to have no significant adverse effects on water quality. The BMPs most likely to be associated with measurable adverse water quality were road stream crossings, developed recreation sites, and water source development. These BMPs also were found to have relatively low effectiveness when implemented.

Effects of Forest Management on Water Supply

The scientific evidence for relations between forests and water supply, however, has been inconclusive (Dudley and Stolton 2003; Troendle et al. 2007). Research has shown that forests have had a limited role in flood protection and variable effects on total water yields and base flows (Ziemer and Lisle 1998; USDA Forest Service 2000; Calder et al. 2007; Moore and Wondzell 2005; National Academy of Sciences 2008). In contrast, several studies have convincingly demonstrated that forests protect water quality by reducing erosion and removing runoff pollutants (e.g., USDA Forest Service 2000; Dudley and Stolton 2003; Calder et al. 2007). Forested watersheds in interior California are the location of California's winter snowpack. In contrast to rainfall that runs off rapidly, these snowpacks store enormous quantities of water through the winter wet season and release this stored water as spring and early summer snowmelt runoff, when it is most needed by humans and the environment, reducing the need for additional downstream dams and reservoirs. (For information on the benefits of increasing snowpack and extending snowmelt via snow fences, refer to resource management strategy report, *Other Resource Management Strategies*.)

Predicted climate changes for California are likely to have large impacts on forest ecosystems and on water supply in the near future. Climate model predictions suggest that there will be a shift in precipitation that results in more rainfall and less snowfall at mid-elevations in the Sierra Nevada (see <http://www.water.ca.gov/climatechange> for more detail), and in fact, more rapid spring snowmelt in the Sierra Nevada is already occurring (Peterson et al. 2008). This predicted shift toward less snow is critically important for water management because the existing water development infrastructure is designed to exploit streamflows that are driven by gradual releases of water during snowmelt. If snow is replaced by rain at mid-elevations, then winter flood peaks are likely to become larger and more frequent, and reservoir storage is likely to be exceeded in wet months when demand is low. Correspondingly, summer stream base flows will be lower in dry months, when demand is high. These climate-driven impacts could

lead to proposals for new dams and reservoirs on forest streams with their resulting environmental impacts, as well as for additional off-site reservoirs.

Climate change also directly affects forests through increased drought stress, which makes trees more vulnerable to insect attack, with the resulting increased rates of tree mortality influencing wildfire frequency, size, and severity. These stresses on forests will affect their capacity to naturally regulate streamflow and buffer water quality. Many streams that are now perennial are likely to become intermittent with the resulting loss of riparian zones, aquatic habitats, and other beneficial uses of water that depend on perennial flows.

The importance of forest management for protection and improvement of water resources has increased due to concerns about increased demand for water, extended drought, economic and environmental costs of new water-supply infrastructure, effects of water transfers on endangered species, and effects of climate change on water supply and hydropower generation. Although current scientific consensus supports the role of forests as protectors of water quality, the potential for improvements in the availability of water through active forest management should not be overlooked (Bales et al. 2011). The following sections discuss forest management actions that have potential for improving water resources in California. (Discussion of an additional management action, snow fences, is in resource management strategy report, *Watershed Management*.) Forest management activities that alter streamflow regimen to benefit downstream water users (primarily by altering the timing of streamflow peaks) may be more successful than attempts to increase total water yield.

Vegetation Management for Water Supply

Management of forest vegetation to improve water supplies has a long history in the western United States. Early efforts attempted to reduce transpiration or increase snowpack by removal of trees, most ending with limited success (Ziemer 1987). Changes in water yields resulting from vegetation management are highly variable and difficult to measure, with indications that treatments must remove at least 20 percent of the vegetation to have a measurable effect on streamflow (Troendle et al. 2007). Computer simulations by Troendle et al. (2007) indicate that every twelve acres of forest thinning (fuels reduction) could theoretically produce an increase of 1 acre-foot (af) of runoff. They suggested that the water yield response to large-scale forest thinning in the northern Sierra Nevada forests would be short-lived with a single treatment, perhaps only 15 years, but that an active management program could result in subtle increases in water yield. Some studies have provided limited evidence that measurable water-yield increases have occurred in larger watersheds in the past in response to vegetation removal. Blanchard (1962, as cited in Zinke 1987) investigated the cumulative effect of 30 years of logging on the South and Middle Forks of the Mokelumne River in the central Sierra Nevada. He reported that between 1930 and 1961, approximately 40,000 acres of forest were logged and that water yields from these watersheds gradually increased during that time period.

Innovative approaches that utilize selective thinning of younger, smaller trees, according to E. Holst of the Environmental Defense Fund, show some promise for limited improvement in streamflow regimen, as well as reducing fuel loading and increasing carbon sequestration (also, Troendle et al. 2007). However, research to date in California indicates a limited potential for increases in water yield following forest vegetation management treatments (e.g., Rector and MacDonald 1987). Most of this research was conducted under conditions of lower stand densities than now exist in California forests. Vegetation

treatments in forests with higher stand densities could produce more benefits to water resources than indicated by research over the past several decades. Research is currently underway to evaluate such benefits (Bales et al. 2011). The effects of vegetation treatments on water availability will likely depend on the extent and intensity of treatments, as well as the length of intervals between treatments. Decisions as to whether to implement specific vegetation management projects will need to consider effects on other forest resources and potential adverse effects to water quality from vegetation treatments. Bales et al. (2011) report that their preliminary estimates based on average climate information suggest that treatments in the Sierra Nevada that would reduce forest cover by 40 percent of maximum levels across a watershed could increase water yields by about 9 percent. These treatments, however, also have potential to increase surface runoff and erosion from disturbed soils (Cram et al. 2007).

Fuels/Fire Management

Wildfire Impacts on Watershed Resources

Wildfires affect water resources by removing vegetation and altering soils and ground cover, with the magnitude of post-wildfire impacts being dependent on burn severity (Ice et al. 2004; Neary et al. 2005; Moody et al. 2008). These changes have large implications to water resources through their effects on transpiration rates, water infiltration rates, rates and magnitudes of erosion, peak and base streamflows, and total water yield.

In the absence of human intervention, wildfires were regular occurrences in California forests, where relatively frequent fires prevented large accumulations of fuel materials and fires were generally fast-moving, low-intensity, and did not kill established trees. Active fire suppression since the 1920s has led to a situation in much of California where forests have developed high fuel loads that greatly increase the risk of catastrophic high-intensity, stand-replacing fires that kill all vegetation, generate large volumes of eroded soil and ash (Robichaud 2000; Reneau et al. 2007; Rulli and Rosso 2007; Carroll et al. 2007), and cause large quantities of mobilized nutrients such as nitrate nitrogen, ammonium nitrogen, and phosphate phosphorus to move into stream runoff (Miller et al. 2006).

The removal of forest canopies that is associated with high burn severity temporarily reduces transpiration and interception losses. Consequently, streamflows increase until vegetative re-growth increases transpiration to or above pre-fire rates (Driscoll et al. 2004), and yields of water from a burned watershed are increased.

In areas with heavy fuels (typically forests and chaparral that have not burned or been treated to reduce fuels for many years) intense wildfire can lead to development of hydrophobic soil layers, particularly in dry coarse-textured soils, that dramatically reduce surface water infiltration rates. The impermeability of hydrophobic soil layers, in conjunction with the lack of ground cover remaining after fires, can lead to increased erosion and early-season surface runoff (Neary et al. 2005; Onda et al. 2007; Moody et al. 2008), causing greater transport of sediment to downstream reservoirs and adverse impacts to water treatment and conveyance facilities (Neary et al. 2005; Moser 2007).

Post-wildfire erosion is highly variable, difficult to predict, and highly dependent on the size, number, and intensity of storm events during the first one-to-two winters following the fire.

Increases in erosion are typically two or more orders of magnitude for intense wildfires the first winter after burning (Robichaud et al. 2010).

Peak streamflows are increased after intense wildfires, but the magnitude of this increase varies greatly by size of the watershed and its location in California. Changes in post-wildfire peak flows are greatest in small watersheds (e.g., < 250 acres) since stormflow response of small basins is controlled primarily by hillslope processes, including infiltration rate, which, in turn, are affected by wildfire (Neary et al. 2005). While data are limited, peak flow increases are likely to be higher in Southern California chaparral-covered basins than in Northern California coastal and snow-dominated watersheds (Robichaud et al. 2000; Neary et al. 2005). Peak flow increases in Southern California are commonly predicted to increase two to three times following intense wildfire for flows that occur with a recurrence interval of two years or greater (Rowe et al. 1949; Moody and Martin 2001).

Although increased water yield is a potential impact of large, intense wildfires, it is generally not significant. Where 75 to 100 percent of the vegetative cover is removed, runoff may increase from 0.1 af per acre burned in watersheds receiving 15 inches of mean annual precipitation, to af per acre burned for watersheds receiving 40 inches of mean annual precipitation (based on Turner 1991). In forested areas, water-yield increases are minimal until basal area loss to fire exceeds 50 percent (Potts et al. 1989).

The additional water yields that result from catastrophic wildfires are generally considered to have little value for water supply and hydroelectric energy generation. Almost all of the additional runoff occurs during the wet season and must be regulated for dry season use by surface reservoir storage (Ziemer 1987). Typically, flows increase during large storm events when water is intentionally allowed to pass through reservoirs owing to flood management concerns. The occasional short-term positive gains from increased water yield are more than offset by the frequent short- and long-term negative impacts of increased peak flows, increased sedimentation, and decreased water quality (California State Board of Forestry 1996).

Increases in suspended sediment and turbidity are usually the greatest impact to water quality following intense wildfire, besides the direct and indirect effects fires can have on water delivery infrastructure. While data are scarce, post-wildfire turbidity values are often expected to exceed drinking water standards for water supplies, and make water treatment more difficult and expensive. Post-fire sediment concentrations are generally highest the first year after the fire, but the extent of sediment mobilization depends on the size of the storms following the fire (Fiori 2005). Increased sedimentation can severely impact aquatic habitat, including that for state and federally listed anadromous salmonids in Northern California. Intense wildfires also remove streamside vegetation, causing water temperatures to rise (Amaranthus et al. 1989; Mahlum et al. 2011). Increased water temperatures can adversely fish species by increasing pathogens and algae, and by decreasing amount of dissolved oxygen and aquatic organisms available to fish (Amaranthus et al. 1989).

Nitrogen is the most important nutrient affected by fire, with the amount of change in nitrogen in a burned area being directly related to the magnitude of soil heating and fire severity, and proportional to the amount of organic matter destroyed (Neary et al. 2005). Intense wildfire can lead to significantly increased nitrogen loads in stream water, particularly in Southern California where post-wildfire concentrations of nitrogen in streams as soluble nitrate have been found to exceed drinking water standards (Meixner and Wohlgemuth 2004), but not in Northern California (Cohen 1982).

Fuel Treatments to Reduce Wildfire Impacts on Watershed Resources

Fuel hazard reduction projects have been shown to reduce the risk of catastrophic crown wildfire (Martinson and Omi 2003; Omi and Martinson 2004), reducing both the severity and frequency of wildfire (Elliot 2010). Fuel reduction projects can have adverse effects on water quality (McClurkin et al. 1987; Wondzell 2001; Grace et al. 2006), but these effects are generally minor and temporary, and are far exceeded by the adverse effects of catastrophic wildfires (Benavides- Solorio and McDonald 2001; USDA Forest Service 2005; Madrid et al. 2006; Hatchett et al. 2006; Cram et al. 2007; Robichaud et al. 2007; Gokbulak et al. 2008). The adverse impacts of wildfire are generally much greater per unit of affected area than the impacts of fuel reduction projects, and also affect much larger areas than are included in fuel reduction treatments.

Prescribed fire, thinning, and mastication are the main types of fuel reduction methods used to decrease the intensity, extent, and negative consequences of wildland fire in California.

Prescribed herbivory (e.g., cattle and goat grazing used to maintain fuel breaks) is an additional option that is sometimes used. The most effective fuel reduction treatments for decreasing the spread and intensity of wildfires have been combinations of mechanical treatments and prescribed burning (Stephens and Moghaddas 2005; Dailey et al. 2008). Fuel management treatments are generally required every 10 to 20 years to maintain their effectiveness in reducing the risk of catastrophic wildfire (Robichaud et al. 2010).

In general, hydrologic impacts from prescribed burning are small, since these fires are usually low intensity (Beschta 1990; Heard 2005; Robichaud et al. 2010). Prescribed burns in chaparral typically generate more soil heating than prescribed burns in either grasslands or forests and produce more sediment than with other vegetation types (10-30 percent of the sediment yields after high severity wildfires) (Wohlgemuth 2001). Prescribed fires in chaparral that kill a significant proportion of the mature canopy or expose more than 35 to 50 percent of the soil can have a significant, detectable effect on annual water yields lasting 8-10 years, but with little detectable impact on downstream water storage reservoirs (Troendle et al. 2010). Nutrient impacts to water quality associated with prescribed burns is minimal in both forested and chaparral watersheds (Stephens et al. 2004; Meixner and Wohlgemuth 2004).

Commercial thinning operations that remove a significant portion of the overstory canopy have the potential to elevate stream sediment loads when the proportion of bare soil is high (Robichaud et al. 2010). Roads associated with commercial thinning operations usually are the largest sediment source associated with commercial timber operations (MacDonald et al. 2006). Only relatively heavy thinning operations can be expected to increase annual water yields in wetter environments, with no measurable increase in runoff expected from thinning operations that remove less than 15 percent of the forest cover or in areas with less than 18 inches of annual precipitation (Reid 2012; Robichaud et al. 2010). Burning of slash piles often produced with thinning produces intense soil heating at the pile locations and alters soil properties, but very limited movement of nutrients downslope from the piles has been detected (Hubbert et al. 2010).

Hydrologic impacts associated with non-commercial fuel reduction thinning operations that are done to reduce the risk of catastrophic wildfire are small, producing only short-lived impacts to runoff and sediment production. Non-commercial thinning to reduce fuel loads is increasingly being accomplished using masticating machines that mechanically grind, crush, shred, chip, and chop fuel. Woody material that remains following mastication increases the amount of ground cover and substantially reduces erosion potential. While research is limited, mastication appears to be an effective thinning treatment for

overstocked timber stands with few negative impacts on soil compaction or soil erosion (Hatchett et al. 2006).

Management Strategies to Reduce Adverse Impacts Associated with Wildfire

Forest management activities to reduce fire severity on California's 18 National Forests are currently administered under the National Fire Plan (NFP) and the Healthy Forest Initiative (HFI). Approximately 70 percent of the 20 million acres of National Forest system lands in California, or 14 million acres, are in need of treatments to reduce fuel loads to natural levels. In all of California, approximately 21 million acres have been designated as high-priority landscape for treatment (California Department of Forestry and Fire Protection 2010a). The USDA Forest Service and other federal and state agencies are currently treating about 220,000 acres per year in California (approximately half with prescribed burning), while an average of 320,000 acres are burned annually by wildfires (California Department of Forestry and Fire Protection 2010a). Prior to European settlement (pre-1800), it has been estimated that 4.5 million acres are burned per year on average in California (Stephens et al. 2007).

Firefighting tactics are increasingly being modified to protect water quality and aquatic organisms (National Wildfire Coordinating Group Training Working Team 2004). Guidelines in effect since 2000 specify that aerial fire retardant drops are to be avoided within 300 feet of waterways (National Interagency Fire Center 2010). Rapid restoration of areas disturbed by fire suppression actions to reduce erosion potential and protect water quality is routinely included in suppression efforts on both National Forest and non-federal lands in California. Fire control lines, particularly those created by heavy equipment, disturb the soil, increase soil compaction, reduce infiltration, can become sources of sediment if not properly rehabilitated, and can alter runoff patterns (Neary et al. 2005; Backer et al. 2004). Practices used to reduce these impacts include installation of proper drainage structures on firelines and roads, and removal of soil from emergency stream crossings built when constructing firelines with crawler tractors.

Following fire containment, burned areas associated with wildfires greater than 500 acres on National Forest lands are assessed, and high-risk areas with downstream values-at-risk are treated to prevent adverse effects on water quality and other resources (Robichaud et al. 2000). Values-at-risk refers to natural resources such as salmonid habitat and human communities that may be adversely affected by the movement of water and sediment from burned areas.

The USDA Forest Service uses its Burned Area Emergency Response (BAER) program to prescribe practices to reduce erosion potential, as well as to reduce threats to life and property. Similarly, at the direction of the governor, California's Emergency Management Agency (Cal EMA), Natural Resources Agency, and Environmental Protection Agency (Cal EPA) assemble multi-disciplinary teams when necessary to assess post-wildfire potential impacts to life and property on state and private lands. Commonly specified measures include notification of residents in areas at risk for debris slides and channel-derived debris flows, use of automated precipitation and stream gauges linked to local government response and flood control agencies for early warning for evacuation, road and stream crossing improvements, installation of structure protection devices (e.g., K-rails), and on USDA Forest Service lands where there are high values-at-risk, such as aerially applied straw mulch, and hydro-mulch (Robichaud et al. 2000; Wohlgemuth et al. 2009). Aerial grass seeding has rarely been used in California after 2000, since it has not been shown to be effective in reducing hillslope erosion and often inhibits native species regeneration (Conard et al. 1995; Wohlgemuth et al. 1998; Beyers 2004). Post-wildfire

assessment programs will likely become increasingly important in the future due to projections of higher frequency and intensity of wildfires related to climate change.

Recommendations

It is recommended that watershed protection be enhanced through the strategic placement of fuel reduction projects in high-priority water supply watersheds, (high-priority water supply watersheds are displayed in Chapter 3 of the *2010 Assessment of California's Forest and Rangelands* [California Department of Forestry and Fire Protection 2010a]) utilizing existing state and federal cost-share programs on non-federal wildlands (California Department of Forestry and Fire Protection 2010b). Fuel reduction projects should use: (1) mechanical thinning treatments that limit ground disturbance, particularly on steeper slopes and more erodible soil types (Cram et al. 2007), and include appropriate road design, construction, and maintenance practices, (2) mastication where slope gradient is appropriate, and (3) low-severity prescribed fire preserving the litter/duff layer and existing nitrogen levels. Fuel reduction treatments, such as thinning, can reduce the threat of high-intensity wildfire, and make California forests more resilient in warmer climates (Bales et al. 2011), as well as providing other ancillary benefits, such as biogeneration of power.

Road Management

Thousands of miles of roads have been constructed through forests in California, primarily to provide access for timber harvest. The 18 National Forests in California alone contain approximately 50,000 miles of forest roads, of which roughly 20,000 miles may no longer be needed for their original purposes (Dombeck 2007). Private forest lands contain many additional thousands of miles of roads. These are mostly unpaved roads and they can have significant effects on hydrology and water quality through their roles in sediment transport and hydrology when they are improperly designed, constructed, or maintained.

Unpaved roads, particularly those adjacent to streams and road stream crossings, are usually the dominant source of management-related sedimentation in forested environments in California due to surface erosion, gullyng, and mass wasting (Cafferata and Munn 2002; USDA Forest Service 2004; MacDonald et al. 2004; Coe 2006; Cafferata et al. 2007b). Excessive sedimentation associated with roads is a concern because of potential negative impacts on stream habitat and water quality from sediment that is discharged either episodically when roads or road-stream crossings catastrophically fail, or chronically from incremental surface erosion. However, a relatively small proportion of the total road length produces most of the road-related sediment delivered to streams (McCashion and Rice 1983; Coe 2006).

Forest roads can have significant effects on hydrology by generating overland flow and intercepting subsurface flow, which increases flood peaks (Jones and Grant 1996) and decreases recession flows. Stream crossings are vulnerable to damage by high flows (Furniss et al. 1998) and can divert streams from their natural channels, resulting in serious erosion and water quality problems (Best et al. 2004).

Roads built to modern standards have reduced impacts to forest streams, but many of the forest roads in California were built decades ago to very low design standards, often in environmentally sensitive locations such as unstable hillslopes and riparian areas. A significant number of older roads are part of the current road network, while others have been neglected and abandoned with no consideration or mitigation of ongoing erosional impacts (Cafferata et al. 2007b). These “legacy” roads are particularly susceptible to catastrophic failure during high magnitude, low frequency storm events, such as the one in 1997 that

caused extensive flooding throughout a large part of Northern and Central California (Furniss et al. 1998; Madej 2001).

Many of these adverse hydrologic and water quality impacts of roads can be reduced by upgrading and replacing culverts, outslipping road treads, and installing road drainage structures such as waterbars and rolling dips at appropriate spacing, particularly near stream crossings (Furniss et al. 1991; Weaver and Hagans 1994; Keller and Sherar 2003). Roads no longer necessary for resource management or recreation can be effectively decommissioned by removal of fills at stream crossings and partial or total outslipping of road treads, including cuts and fills (Madej 2001; Cook and Dresser 2007). Road decommissioning can potentially reduce water quality impacts. However, it can be difficult to find roads producing significant impacts that people agree should be decommissioned.

Detailed field surveys are the main tool available to identify the road segments of greatest concern (Weaver et al. 2006; Korte and MacDonald 2007). Public and private landowners in California are actively inventorying their road networks, prioritizing road segments requiring road improvement or decommissioning work, and completing projects. A considerable amount of road upgrade work has been completed to date with both public and private financing.

While there are short-term impacts associated with road improvement and decommissioning, particularly at stream crossings, improved operator practices has lessened these effects (Pacific Watershed Associates 2005; Cafferata et al. 2007b), and treatments will reduce the long-term sediment production overall from older roads (Madej 2001).

Riparian Forests

Riparian forests are forested lands, usually in narrow linear strips, that are located immediately adjacent to streams, lakes, or other water bodies. These communities occupy a transition zone between aquatic and terrestrial habitats, and are distributed in complex patterns that are responses to geomorphology, annual flood timing and extent, soil moisture, and plant competition. The boundaries between riparian and upland forests are not always distinct, and the width of a riparian forest strip varies laterally throughout the channel network and is strongly influenced by geomorphology (Naiman et al. 1998).

In recognition of the central role played by riparian forests in the landscape, the California Forest Practice Rules require that the beneficial functions of riparian zones and populations of native aquatic and riparian-associated species must be maintained where they are in good condition, protected where they are threatened, and restored where they are impaired insofar as is feasible. Unfortunately, riparian forests are prone to invasion by noxious non-native plant species that reduce the value of the riparian community to humans and wildlife.

Forested floodplains are zones of very high biological diversity, generally harboring the highest biodiversity of both terrestrial and aquatic organisms within the watershed landscape (Naiman et al. 1998) and providing important habitat for wildlife (Kattelman and Embury 1996; Ligon et al. 1999). Riparian floodplains play large roles in forested watersheds that are disproportional to their small area in the landscape. The high surface roughness of forested floodplains has large effects on stream hydrology by reducing floodwater velocities and spreading flood flows across a larger area of the floodplain. The

retention and slowing of floodwaters across a wider area allows floodwaters to recharge alluvial groundwater aquifers and attenuates downstream flood flows (Cafferata et al. 2005).

Studies have shown that riparian forests can improve water quality. Riparian forests contribute to reductions in sediment, nutrient, and pesticide loads of surface runoff through physical and biological processes, reducing these inputs to watercourses. Canopy shading by riparian trees reduces stream water temperatures, which is important for many fish species that are adversely affected by elevated water temperatures. In cooler coastal areas, canopy shading may restrict primary productivity (Wilzbach et al. 2005).

Riparian forests are protected on federal, state, and private timberlands by regulating areas near streams as riparian buffers, within which management actions such as timber harvesting and road building are regulated. The width of riparian buffers, and restrictions on management activities within them, are based largely on land ownership. Within the National Forests, riparian buffer widths vary based on planning province standards and guidelines, with riparian protection being most extensive for the six National Forests that operate under the Northwest Forest Plan. Provincial refers to the three major planning provinces used in the National Forest System in California — the Northwest Forest Plan province, the Sierra Nevada Framework province, and the Southern California province.

Even with these protections, the extent of riparian forest is greatly diminished from its historical extent, particularly in lowland valleys where riparian forests have been converted to orchards and other agricultural uses. In the Central Valley, riparian forests historically covered more than 900,000 acres but presently account for less than 100,000 acres (Barbour et al. 1993).

Unmanaged riparian stands can be sources of rapid fire migration in fire-prone landscapes (Murphy et al. 2007). Fuels reduction within riparian buffers may be needed in some cases to reduce threats of catastrophic wildfires, particularly in the interior parts of California (USDA Forest Service 2007; Van de Water and North 2011). Goals for this type of work include creating fire resilient forests, promoting reduced fire intensities, and retaining functional aquatic and riparian habitat following a wildfire. Removal of trees from riparian buffers remains highly controversial (Welsh 2011), and forest management and regulatory agencies are carefully evaluating monitoring data, particularly with regard to the use of mechanical equipment in streamside zones (Norman et al. 2008).

Some riparian forests are used for livestock grazing, usually within allotments that consist mostly of upland pasture. The availability of water and forage make riparian areas attractive to livestock, which can damage riparian forests through trampling, browsing, and contamination of streams with fecal material (Campbell and Allen-Diaz 1997). BMPs for range management and National Forest standards and guidelines for riparian management are designed to protect riparian forests from damage by livestock. Although exclusion of cattle may be needed during and immediately after restoration of riparian forests, grazing strategies that minimize impacts on riparian forests through restrictions on livestock numbers, distribution, and season of use can be used to eliminate the need for permanent fencing.

Riparian forests are primarily a result of the interplay of hydrologic processes occurring in the stream, floodplain, and groundwater, and alteration of these processes can have dramatic effects on riparian plant communities. Many of the tree and shrub species that dominate California riparian forests (e.g., cottonwood, willow) are dependent for regeneration on the annual flooding and exposure of a floodplain

because their seeds can only germinate in moist, emergent areas that have been scoured of vegetation – conditions that occur only after high water events.

Dominant riparian tree species often have little or no tolerance for dry conditions and require the reliable source of moisture supplied by the streamflow. Some riparian plant species have shallow root systems and can only utilize water in shallow areas of the soil profile or in the stream channel directly, while other species have roots that penetrate deeper into the soil profile and utilize available groundwater, which is typically replenished by streamflow infiltration.

Species distributions are further affected by patterns of sediment deposition caused by stream hydrologic processes. Riparian plants, particularly trees, also affect stream hydrology, in turn, by contributing large woody debris that creates pools, and affect channel morphology through the actions of roots.

Riparian forests, therefore, may affect and be effected by channel incision and groundwater storage in much the same ways as meadows. The water quantity and quality benefits provided by riparian forests can be preserved and enhanced through actions that maintain natural channel geomorphology. Thus, protection of riparian forests depends heavily on effective management of upland watersheds to prevent excessive runoff and sedimentation, as well as control of non-native invasive species.

Meadow Restoration and Groundwater

Meadow wetlands are level or gently sloping alluvial landforms with seasonally high water tables that support flood-tolerant herbaceous plants (Ratliff, 1985; Weixelman et al. 2011). Geologic and climatic factors have resulted in the formation of numerous meadows in the mountains of California, with most meadows located in the Sierra Nevada. These meadows, in aggregate, comprise a small part of the landscape, but they provide disproportionately important ecological services owing to their role in retaining water at or near the land surface during California's long dry summers.

In their natural condition, meadows have shallow meandering channels that allow high flows to spread across the meadow surfaces, depositing suspended sediments and recharging meadow aquifers (Wood 1975). These meadows act as natural reservoirs, storing and releasing snowmelt and rainfall runoff that passes over and through fine-grained, sod-covered meadow deposits (Loheide and Gorelick 2007; Cornwell and Brown 2008; Hammersmark et al. 2008; Tague et al. 2008; Ohara et al. 2013).

By 1940, many meadows throughout the Sierra Nevada had been eroded by incised channels, or gullies, as a result of unrestricted livestock grazing, road building, railroad construction, elimination of beavers, and other causes (Hughes 1934; Kraebel and Pillsbury 1934; Wood 1975; Ratliff 1985; Kattelman and Embury 1996; Martin 2006). A recent survey found that more than half of Sierra Nevada meadows are eroded by incised channels (Fryoff-Hung and Viers 2013). Current activities are designed to avoid damage to meadows, but the effects of earlier practices remain on the landscape and are unlikely to heal without active restoration (Ratliff 1985). Future disturbances from wildfires, intense storms, encroachment by conifers, and illegal activities could cause further damage that will require restoration.

Eroded meadows lose their capacity to store groundwater (Loheide and Gorelick 2007; Cornwell and Brown 2008; Hammersmark et al. 2008; Tague et al. 2008) because deeply incised channels can convey flood peaks without overbank flooding and recharge, and because groundwater is drained more rapidly

from meadow aquifers. In addition, channel erosion in meadows adds to stream sediment loads through bank erosion and headcut retreat, adversely affecting downstream water quality and reservoir capacity (Micheli and Kirchner 2002). Loss of groundwater to evapotranspiration, however, is often reduced by meadow erosion and increased by restoration (Loheide and Gorelick 2005).

In the 2009 State Water Project (SWP) update, potential groundwater storage benefits of meadow restoration were estimated at 50,000 to 500,000 acre-feet (af) annually. The large uncertainty in these estimates resulted from the limited information available at that time regarding meadow areas, erosion depths, and hydraulic properties.

Based on preliminary results of an ongoing regional assessment (Fryoff-Hung and Viers 2013), the potential groundwater storage benefits of meadow restoration throughout the Sierra Nevada are likely to range from 25,000 to 50,000 af annually. These estimates are lower than the estimated range in the 2009 SWP update because recently acquired measurements indicate that meadow areas and erosion depths are lower than initially estimated for the 2009 SWP update.

The reduced soil moisture and elimination of surface flooding that is associated with meadow incision leads to changes in the associated plant community to types that have less value and provide fewer ecological services. Drying of meadow soils allows invasion by drought-tolerant brush and conifer species that contribute to heavy fuel loading and add to the risk of catastrophic wildfires (Allen-Diaz 1991; Dwire et al. 2006; Berlow et al. 2002; Loheide and Gorelick 2007). Loss of high quality forage provided by wet-meadow sedges, rushes, and grasses decreases forage value for meadows that are grazed by livestock (Ratliff 1985; Stohlgren et al. 1989). Loss of wetland vegetation reduces habitat area for several rare, threatened or endangered species such as the Mountain Yellow-legged Frog, Willow Flycatcher, and Bell's Vireo.

When groundwater is available, but not so persistent as to induce anoxia or root rot, woody species such as lodgepole pine and sagebrush may transpire groundwater at rates similar to or possibly higher than wet meadow herbaceous vegetation (e.g., Lautz 2007; Tague 2009). Woody plants at some elevations continue to transpire throughout the year (although at lower rates during the winter; Bales et al. 2011), adding to transpiration losses, whereas herbaceous meadow plants senesce in the fall and do not transpire for about half of each year (e.g., Wood 1975). Removing encroaching conifers or sagebrush may in some cases result in reduced groundwater transpiration losses (Cafferata 2005). This might extend the period that meadow areas are wet longer into the summer season, but it would not be expected to produce a major shift in groundwater trends. The effect on summer groundwater elevation would decrease over time if coniferous or shrub vegetation was allowed to regenerate.

The USDA Forest Service and other agencies and organizations have been restoring meadows for many decades (Kraebel and Pillsbury 1934; Hughes 1934). In the past 20 years, the pace and scale of meadow restoration have accelerated owing to increased external support and the advent of the "plug and pond" restoration methodology. This acceleration of meadow restoration has raised concerns among some downstream water users that restoration may decrease summer streamflows, and has highlighted the need to better understand meadow hydrology and the manner in which restoration affects downstream flows.

The effects of meadow restoration on streamflow are not well documented except in a few cases. Streamflow records in the Feather River watershed do not show clear evidence for increased flows

downstream of restoration projects, and in fact show at least temporary decreases in some restored meadows (Hoffman et al. 2013). However, recent scientific publications have generally indicated that meadow restoration will benefit downstream flows during dry periods (Tague et al. 2008; Hammersmark et al. 2008; Ohara et al. 2013). Long-term effects of restoration on downstream flows are likely to vary depending on climate, geology, vegetation, and land use.

Urban Forestry

Trees planted along streets and in city parks, lots, and private residences collectively form urban forests, and urban forestry practices address the maintenance of existing urban trees as well as the planting of new trees in and around cities. Although urban forests are not managed specifically for natural resource production or conservation, they have environmental benefits that extend well beyond aesthetics.

Urban areas in California cover roughly 5 percent (7,944 square miles) of the land base, but support 94 percent of the population (California Department of Forestry and Fire Protection 2010a). An estimated 15.1 percent of California's urban area (800,000 acres), which is home to almost one-third of the state's population (9.5 million people), is associated with high threats from air pollution and urban heat islands (California Department of Forestry and Fire Protection 2010a). Urban trees are an important means of mitigating heat and air pollution. As a result, communities throughout California are recognizing the importance of urban trees and have plans to expand urban forests. The need for expanding or enhancing urban forests is substantial, with 372 communities identified as high priority for tree planting in urban areas by the *2010 CAL FIRE Forest and Range Assessment* (California Department of Forestry and Fire Protection 2010a).

Urban Watershed Forestry

While not part of the wild environment, urban trees contribute to the overall health of a watershed, and their contribution is addressed by the discipline of urban watershed forestry, which is an integration of urban and community forestry and watershed planning. Urban and community forestry focuses on how to manage urban forests for environmental, community, and economic benefits while watershed planning focuses on strategic land use and resource management within a watershed. The integration of these two methods into urban watershed planning recognizes the role trees play in protecting water resources, and is becoming a valuable resource management tool for urban planners.

Tree Cover and Watershed Benefits

Trees in an urban setting provide multiple watershed benefits (Table 2), including reduction of stormwater runoff and stream channel erosion, improved soil and water quality, and reduction of air and water temperatures. For example, it is possible for a single tree to contain 100 gallons of water or more within its leaves and bark, which when multiplied by the many trees in an urban setting, produces an impressive retention capacity that can reduce stormwater runoff by 2-7 percent (Tree City USA 2010). In conjunction with other landscaping, an estimated 65 percent runoff reduction can be achieved (Tree City USA 2010), and water retention systems such as vegetation swales, stormwater basins, structural soils, tree pits, and riparian buffers improve runoff reduction even more.

Stormwater Runoff

Trees reduce stormwater runoff by using soil water through transpiration and intercepting rainwater on leaves, branches, and tree trunks, which changes runoff quantity and pollutant loads in several ways. rates,

Table 2 Watershed Benefits of Urban Forest Cover

Benefit	Description
Reduce storm water runoff and flooding	<ul style="list-style-type: none"> • Trees intercept rainfall in their canopy, reducing the amount of rain that reaches the ground. A portion of this captured rainwater evaporates from tree surfaces. • Trees take up water from the soil through their roots, which increases soil water storage potential and lengthens the amount of time before rainfall becomes runoff. • Trees promote infiltration by slowing down runoff and by increasing soil drainage in the root zone. The addition of organic matter (e.g., leaves) also increases storage of water in the soil, further reducing runoff. • Forested land produces very little runoff, which can reduce downstream flood flows that erode stream channels, damage property, and destroy habitat.
Improve regional air quality	<ul style="list-style-type: none"> • Trees absorb pollutants such as nitrogen dioxide, carbon monoxide, ozone, and particulate matter from the atmosphere. • Trees reduce air temperature, which reduces formation of pollutants that are temperature dependent, such as ozone. • Trees indirectly improve air quality by cooling the air, storing carbon, and reducing energy use, which reduces power plant emissions.
Reduce stream channel erosion	<ul style="list-style-type: none"> • Trees growing along a streambank prevent erosion by stabilizing the soil with root systems and the addition of organic matter.
Improve soil and water quality	<ul style="list-style-type: none"> • Trees prevent erosion of sediment by stabilizing the soil, and by substantially dispersing raindrop energy. • Trees take up stormwater pollutants, such as nitrogen, from soil and groundwater. • Forested areas can filter sediment and associated pollutants from runoff. • Certain tree species break down pollutants commonly found in urban soils, groundwater, and runoff, such as metals, pesticides, and solvents.
Provide habitat for terrestrial and aquatic wildlife	<ul style="list-style-type: none"> • Forests (and even single trees) provide habitat for wildlife in the form of food supply, interior breeding areas, and migratory corridors. • Streamside forests provide habitat in the form of leaf litter and large woody debris for fish and other aquatic species. • Forest litter such as branches, leaves, fruits, and flowers form the basis of the food web for stream organisms.
Reduce summer air and water temperatures	<ul style="list-style-type: none"> • Riparian forests shade the stream and regulate summer air and water temperatures, which is critical for many aquatic species. • Trees and forests shade impervious surfaces, reducing temperature of stormwater runoff, which can minimize the thermal shocks normally transmitted to receiving waters during storms.

Source: Watershed Forestry Resource Guide - Urban Watershed Forestry 2008

and interception of rainfall by the canopy reduces the volume and timing of runoff and reduces soil erosion caused by impacts from raindrops.

Structural Soils

Urban areas are challenged by extensive impervious surfaces, damaged soils, and little room for greenspace or for stormwater management facilities. In 2004, a collaboration of researchers from Virginia Polytechnic Institute, Cornell University, and University of California, Davis formed a work group to study the use of trees and structural soils to improve water quality. The system developed and evaluated by the group utilized stormwater BMPs to reduce peak flow, reduce runoff volume, and remove pollutants. The system works by guiding the water into a structural soil retention area beneath the pavement where it is absorbed by soil infiltration and root uptake for tree transpiration. Trees have the potential to develop full canopies that result in increased water interception because the reservoir offers a large root area. Tree roots take up excess nutrients and water in the soil reservoir and can enhance infiltration into the subsoil. Together, trees and structural soils can create a zero runoff site. The group found that with such a system it was possible to distribute stormwater management by taking advantage of the mitigation services provided by urban trees (Xiao and McPherson 2009). It also created an alternative to detention ponds in urbanized areas.

Quantifying Benefits

Urban trees have multiple co-benefits. A large deciduous canopy tree can intercept 760 gallons of rainfall in its crown annually and aid in reducing runoff of polluted stormwater and flooding, a benefit valued at \$6 annually on the basis of local expenditures for water quality management and flood control (USDA Forest Service 1999). Larger potential for canopy interception increases the beneficial effects of tree interception of rainfall, with these effects being greater in larger trees and evergreen trees. An evergreen camphor tree, for instance, is estimated to intercept 4,000 gallons annually, providing even greater benefits than a deciduous tree of similar size (USDA Forest Service 1999). In addition, shade from urban forests reduces energy use of city residents by reducing temperatures inside buildings and lowering energy usage rates for interior cooling.

Urban trees also offset greenhouse gas emissions and provide larger-scale climate benefits through their persistent sequestration of carbon in woody material. A study in San Francisco found that urban trees within the city annually sequester an estimated 2,271 tons of carbon dioxide (CO₂) indirectly reduce energy plant emissions by 257 tons of CO₂, representing an estimated value of \$2.3 million annually (Maco et al. 2003). The combined value of this benefit (e.g., carbon sequestration and offset from reduction in energy use) was estimated at \$37,907 annually (Maco et al. 2003). Considering that San Francisco has a mild climate with cool summers, the benefit can be substantially higher in warmer inland cities. There are an estimated 188.5 million urban trees statewide that sequester approximately 414,000 metric tons of carbon annually (or approximately 1.52 million metric tons CO₂) (Novak et al. 2009), so the contribution of urban trees to atmospheric carbon dynamics is substantial.

Recommendations

- Fund urban tree planting in high-priority communities, which should yield multiple water use benefits, such as reductions in stormwater runoff and improved water quality, among other benefits such as air pollution mitigation and reduced energy use. The 2010 Forest and Range Assessment (California Department of Forestry and Fire Protection 2010a) identified 372 communities as high-priority areas for urban tree planting in order to conserve energy or improve air quality.

- Preserve space for large-statured trees in new developments and create such space in developed areas that currently do not have adequate planting sites. Preserving and planting large-statured trees will have a large beneficial impact and improve the extent of urban tree canopy in priority areas. Additionally, improved management of existing urban forest resources will assist in maximizing the benefits of current tree canopy while minimizing long-term costs.
- Encourage and implement BMPs that promote urban forestry for urban stormwater management, which take advantage of benefits offered from tree canopy interception for reduced peak stormwater flows, reduced runoff volume, and removal of pollutants. Use of a variety of stormwater management techniques should be encouraged to maximize urban tree benefits to water resources.

Forest Loss and Fragmentation

The subdivision of large forest parcels and their subsequent development for rural residential uses can cause impacts on water quantity and quality. During the construction phase, there can be increased erosion and sediment delivery to streams. Over the long term, roads and increased impervious surface may change the timing and magnitude of peak streamflow events in small watersheds. Roads and stream crossings also are the main sources for long-term increases in sediment delivery. Although studies comparing the impacts of forest management versus rural residential land uses are rare, there is evidence that rural residential roads produce more sediment than logging roads. Rural residential uses also may affect water quantity at the small watershed scale. These uses will often rely on springs and diversions for water supply. The result may be reduced summertime base flows in streams with attendant biological consequences. The regulatory framework controlling forest management uses is much more demanding than the controls on rural residential uses. In particular, the California Forest Practice Rules require practices to control erosion and sediment delivery. In contrast, most rural counties do not have grading ordinances or similar regulations, especially in relation to private roads (Harris and Cafferata 2005).

Recommendation

Encourage the use of conservation easements to permanently conserve California's private forestlands. Conservation easements are a cost-effective and permanent method of conserving vital water infrastructure, and they could be used more extensively to promote long-term forest stewardship in private forestlands, particularly where land conversion projects are common. The goal is to secure proper watershed functions through conservation easements that permanently conserve forest landscapes, avoiding conversion to housing subdivisions and agricultural lands.

Illegal Marijuana Cultivation

In the past five years, increased impacts from commercial-scale illegal cannabis growing operations have been documented in forested counties throughout California and particularly in the California coast ranges, both on public and private lands. While largely anecdotal, without specific data on numbers of watercourses affected, the impacts have been well documented with digital photographs taken during law enforcement operations (Giusti 2012).

Illegal growing activities adversely impact watershed resources in three main ways: (1) illegal diversions of water from tributary streams utilize low summer flows required for sustaining state and federally listed anadromous salmonids and other species; (2) illegal grading and road building operations cause surface

erosion and slope instability, which produces accelerated sedimentation; and (3) large-scale use of pesticides, fertilizers, and rodenticides adversely impacts water quality.

Typical commercial-scale illegal marijuana gardens found on public land include approximately 7,000 plants, with each large plant using approximately one gallon of water per day (Mallery 2011). This equates to approximately 7,000 gallons of water per day over a period of 3 to 4 months or about 2 to 2.5 af per year per commercial-scale operation. The 2010 Mendocino County Grand Jury Report estimates that only 10 percent of illegally grown marijuana is confiscated annually. More than 500,000 plants are confiscated in many years. Assuming those years are representative, an estimated 5,000,000 plants are produced annually, according to G. Giusti of the University of California Cooperative Extension, Ukiah.

The greatest impact to water resources by illegal marijuana plantations is often not the absolute size of the diversion, but the size of the diversion in relation to the stream being diverted in that it is not unusual for all of the streamflow from a watercourse to be illegally diverted for irrigation using dams, pumps, and elaborate water distribution systems (Thorsen 2011; Mallery 2011). Use of large off-site water storage devices, such as 50,000 gallon water “bladders”, has also been documented. Water diversion causes early de-watering of intermittent streams during a critical time of year for juvenile fish.

Illegal and unregulated grading operations, including grading that enables marijuana cultivation, have been documented in several North Coast counties and they have been found to have adverse watershed impacts. Illegal grading operations increases suspended sediment concentrations and turbidity in intermittent and perennial fish-bearing watercourses, adversely impacting both macroinvertebrates and anadromous salmonids. The extent of illegal grading and its significance to anadromous fish species is currently unknown, however, since counties lack adequate staff to monitor for illegal and/or improper grading (Harris 2011).

Unregulated pesticide, fertilizer, and rodenticide use is extensive in commercial-scale operations and potentially presents a major problem for water quality (Giusti 2012). Mallery (2011) states that an estimated 1.5 pounds of fertilizer are used for every 10 marijuana plants. A three week- long, multi-agency law enforcement operation on public and private lands in Colusa, Glenn, Lake, Mendocino, Tehama, and Trinity Counties in 2011 removed 5,459 pounds of fertilizer and 149 pounds of pesticides from cultivation sites (USDA Forest Service 2011c). An average 5 acre site can contain 20 pounds of rodenticides, 30 bags of fertilizer, plant growth hormones, insecticides, herbicides, and fungicides as well as other chemicals (Mallery 2011). Unused or abandoned chemicals are typically left on-site and leach into waterways and groundwater aquifers, and gasoline and other petroleum products also produce water quality impacts (Giusti 2012).

While the total number and area covered by illegal commercial-scale marijuana operations is unknown, the fact that they operate outside of laws and regulations governing water diversion and water quality protection indicates that they are producing significant impacts wherever they occur. The North Coast Regional Water Quality Control Board is proceeding with an effort to regulate marijuana cultivation on private lands in conformance with the state’s non-point source policy by addressing potential water quality impacts resulting from grading activities, road construction, road maintenance, and hazardous materials handling.

Climate Change

Forests will play an increasingly important role in protecting California's watersheds and associated water supply as the climate warms and precipitation patterns become increasingly variable. Climate change impacts on California's forests that have been measured in the past 100 years include a 10 percent decrease in snowpack, changes in streamflow timing, increased wildfires, and more severe pest outbreaks (California Department of Water Resources 2008).

While susceptible to anticipated changes, proper management of forest habitat provides both climate change adaptation and mitigation benefits. The USDA Forest Service has prepared a resource titled *Responding to Climate Change in National Forests: A Guidebook for Developing Adaptation Options* (USDA Forest Service 2011a). The guidebook is based on the "science-based principles, processes, and tools necessary to assist with developing adaptation options for national forest lands," which can be useful for all forest managers seeking guidance on climate change. One of the key components of successful adaptation in forests will be long-term monitoring and research on the various recommendations and policies that are currently promoted and an adaptive management approach that allows incorporation of new information into the existing management paradigm.

Adaptation

Many existing forest management practices can promote resilience to climate change, and in fact, the best way to ensure successful implementation of high-priority actions is to integrate climate adaptation into existing planning and operational processes. Strategic forest road management will be important in areas prone to flooding and erosion, which can significantly affect water quality due to sediment transport. Incorporating anticipated climate change impacts and vulnerabilities into road management plans and policies will ensure that priorities are based on the changed conditions under which forest roads will need to be managed in the future. Fuel reduction plans should also incorporate climate change considerations so that the threat of high-intensity wildfire situations can be reduced.

Restoration, protection, and proper management of meadows can provide increased water storage and flood protection benefits, which will be very important since increasingly extreme storm events are an anticipated impact of climate change, and precipitation is expected to fall more frequently as rain rather than as snow at lower elevations (California Department of Water Resources 2008). Protection and restoration of headwater streams, including conifer growth in the riparian corridor, could buffer against increasing stream temperature as well as provide habitat connectivity. Healthy forests protect biodiversity, which will be an important buffer against climate change impacts (USDA Forest Service 2011b).

Mitigation

California's forests are carbon sinks, and thus are an important part of climate change mitigation. Sustainable forestry management practices that protect ecosystem services provide greenhouse gas reduction through carbon sequestration as well as other benefits such as water quality protection and energy savings.

Fuel reduction projects, such as mechanical thinning and low severity prescribed fires, initially entail GHG emissions, but could reduce the threat of high-intensity wildfire and thus prevent even greater emissions at a future date as well protect carbon sequestration capacity of remaining trees in thinned

stands. Likewise, managing forest roads uses energy in the short-term, but could result in overall energy savings through reduced sediment transport during heavy rainfall events, thus reducing the energy needed to treat the water downstream.

Urban forestry provides multiple benefits related to climate change mitigation such as decreasing and filtering stormwater runoff, reducing ambient summer air and water temperatures, and carbon sequestration. Careful maintenance of existing urban trees may help offset the “urban heat island” effect and reduce the amount of energy used for cooling in the summer months.

Potential Costs

Vegetation Management

Unit costs for vegetation management on private forest lands in California vary between \$20 and \$1,200 per acre, depending on the methods used. Manual removal of undesirable species ranges from \$70 to \$1,200 per acre. Herbicide applications range from \$20 to \$250 per acre. Herbivory costs range from \$500 to \$1,200 per acre. Mechanical treatments cost between \$800 and \$1,200 per acre. M. Land, of the USDA Forest Service, reported in 2008 that unit costs for vegetation management on National Forest System lands in California are generally higher, ranging from approximately \$1,000 to \$2,000 per acre. On federal land, some or all of the costs of vegetation management can be offset by revenues from commercial timber sales.

Fuels/Fire Management

Unit costs for prescribed fire on private forest lands in California are up to \$500 per acre for grass and shrub fuels and higher for heavier fuels. R. Griffith, of the USDA Forest Service, has stated that unit costs for fuel reduction projects on National Forest System lands in California ranged from \$144 to \$2,476 per acre between 2004 and 2006, with an average unit cost of \$593 per acre.

Road Management

Road upgrading or “storm-proofing” is used to reduce the potential for sediment delivery to stream channels for roads that will remain in service. Recent unit cost estimates for storm-proofing roads on National Forest System lands in the Coast Ranges ranged from \$6,520 to \$13,580 per mile. Road decommissioning is generally much more expensive due to greater planning, heavy equipment use, and hauling costs.

Riparian Forests

No unit cost information is available for riparian forest protection, improvement, or restoration. Actions to benefit riparian forests include appropriate management in both the riparian zone (Van de Water and North 2011; Liquori et al. 2012) and upland watershed improvement projects. Unit costs for upland projects should reasonably represent unit costs for riparian forests.

Meadow Groundwater Storage

Costs of recent meadow restoration projects, including planning and environmental compliance, range from approximately \$1,000 to \$2,500 per acre, according to hydrologist Craig Oehrli of the USDA Forest Service and soil scientist Randy Westmoreland of the Tahoe National Forest, with the higher costs being

associated with projects that require construction of new channels using heavy equipment and end-hauled materials. Maintenance costs for meadow restoration projects are generally very low.

Urban Forestry

The costs of urban tree planting and maintenance can vary greatly with location, site conditions, and the type of tree planted. Total planting cost in California can vary between \$45 and \$160 per tree. After trees are established, maintenance costs are initially minimal, but begin to accrue after about 10 years when trees start to require pruning and hardscape damage from roots needs to be repaired. These maintenance costs can be reduced by careful selection of trees and planting sites. Additional maintenance costs include inspection, administration, legal claims, disease control, removals, and storm litter cleanup. Maintenance costs are typically higher for trees planted in public spaces, since they require more frequent pruning to avoid interference with power and telecommunications lines, and are also generally adjacent to streets and sidewalks. Average annual tree maintenance costs in California, including planting and maintenance, vary from \$13 to \$65 annually per tree, with costs higher on public versus private lands (McPherson et al. 2005).

Illegal Marijuana Cultivation

Mallery (2011) states that the average cost of cleanup for a 10-acre site on public lands involving the use of a helicopter is \$5,000. When environmental remediation is included, the cost of site processing doubles, bringing the average to approximately \$10,000 per 10 acres. These expenses include helicopter fees, fuel consumption, wages, food, gear (tents, hard hats, gloves, shovels, etc.), trash disposal fees, and other variable costs not including the cost of raids, eradication, or investigations (Mallery 2011). Removal of water storage detention basins requires care to restore original flow patterns, while minimizing sedimentation and changes to perennial and intermittent streamflows. Additionally, the removal of miles of irrigation tubing is one of the most intensive parts of remediation efforts, in terms of time, effort, and cost (Mallery 2011). Agencies such as the USDA Forest Service have to divert funding from their primary land management functions to finance cleanup efforts because of the high cost cleaning up illegal sites. Law enforcement incurs added costs when needed to address illegal operators before environmental cleanup efforts can take place.

Major Implementation Issues

The issues described in this section are challenges for implementing one or more of the activities described in the “Quantifying Benefits” section.

Information Needs

Forest management agencies and private timber companies are conducting a number of long-term studies in forested watersheds, including Redwood Creek, Caspar Creek, and South Fork Wages Creek in the northern part of the Coast Ranges; Little Creek in the central part of the Coast Ranges; Judd Creek and Battle Creek in the northern Sierra Nevada; Frasier Peak Creek and Bear Trap Creek in the central Sierra Nevada; and Speckerman Creek, Big Sandy Creek, and the Kings River Experimental Watershed in the southern Sierra Nevada. These studies are providing valuable information about the effects of forest management activities on water quality and quantity, particularly related to timber harvesting, road building, and fuel treatments.

Continued monitoring and additional studies are needed to better understand the effects of forest management activities on water quantity and quality over the wide range of climatic and physiographic conditions found in California. The processes and pathways by which water arrives at the land surface as rain or snow and then reaches stream channels, profoundly affects streamflow regimen, erosion, and contaminant transfer, but these processes are generally poorly understood. Methods for estimating evapotranspiration from different vegetation types need refinement and field verification. Knowledge of groundwater recharge, flowpaths, and storage is limited for mountainous forested watersheds, especially those underlain by fractured rocks. Sources of sediment, transport mechanisms, and the relative importance of erosional processes are not well documented.

Monitoring of streamflow to detect effects of land use is most useful on headwater streams that are not affected by artificial regulation or diversion (MacDonald and Coe 2007). A statewide network of 886 streamflow monitoring stations is operated in California by the U.S. Geological Survey (USGS), but only 214 of these gauges are on streams with more than 50 percent forest cover. Only 31 of these are long-term stations (20 or more years of record) on unregulated and undiverted streams, according to C. Parrat of the USGS, and very few of these stations include water quality monitoring. That density is an average of one long-term stream gauge on an unregulated and undiverted stream for every 1,893 square miles of forest in the state, and some of these stations are in danger of closure due to inadequate long-term funding. A higher density of stream gauges and water quality monitoring stations would be helpful for understanding the distribution, timing, and quality of streamflow from forested watersheds across the state.

Coordination Needs

Forest owners and management agencies have disparate management objectives and constraints, and forest ownership boundaries rarely coincide with natural watershed boundaries, which lead to fragmented, uncoordinated activities that are potentially not effective over the entire watershed. For example, USDA Forest Service funds and staff can generally be used only for work on National Forest System lands, state agencies are frequently prohibited from working on federal lands, and many watershed improvement grant programs are limited to non-federal agencies and organizations. Increased coordination among state, federal, and tribal, private, and non-profit forestry and watershed agencies would provide better opportunities to increase protection of water quality.

A prime example of successful coordination was announced on August 28, 2012 when the Karuk Tribe and the USDA Forest Service signed a Memorandum of Understanding that will protect the Katimiin Cultural Management Area, near present-day Somes Bar, California. This agreement will restore a sacred landscape by using both Karuk traditional knowledge and management practices and the Klamath National Forest Land and Resource Management Plan, which is administered by the Six Rivers National Forest. The agreement will help prevent wildfires as well as build an understanding and cooperation between cultures, the people, and the environment.

Limited Funding for Forest Watershed Restoration

The rate of progress of meadow restoration, road storm-proofing and decommissioning, and vegetation treatment work is largely limited by available funds. In recent years, appropriated federal funding for watershed programs on National Forests has decreased, and revenue-generating timber sales have declined since the mid-1980s. A large proportion of funding for watershed restoration and fuel treatments

is now supplied through state bond measures and grant programs. Some grant programs, however, require non-federal matching funds, which limits the eligibility of projects on federal forest lands.

New sources of funding are needed to continue making progress in watershed restoration. Management of forest resources often results in benefits in water supply, flood control, and flow regulation to downstream communities, whose residents, in most cases, are unaware of these benefits of upstream forest management. With an appropriate outreach effort, these communities, which do not usually contribute to the funding of upstream forest management, might be willing to contribute if the costs and benefits could be demonstrated to them.

Regulatory Requirements

Forest management actions that affect the amounts or timing of streamflow, such as attenuating flood peaks and increasing infiltration to groundwater, may be viewed as threats to existing appropriated rights, and could potentially result in water rights litigation. Surface waters may be appropriated by landowners and other users and these appropriators have legal rights to the water they are permitted to divert. In most cases, reduced flood peaks will not result in less available water for downstream users, but water rights may need to be resolved for any additional water made available by meadow restoration, vegetation management, and fuels treatment.

Harvesting of timber on non-federal lands must comply with the California Forest Practices Act and Rules, the California Environmental Quality Act (CEQA), and other state regulations. Timber harvests and other vegetation and fuel management projects on federal lands are analyzed following NEPA guidelines, and appropriate BMPs are determined for protection of water quality. Federal and non-federal timber harvests, vegetation management, and fuels projects are also regulated by the RWQCBs through Waste Discharge Requirements (WDR) or Waivers of Waste Discharge Requirements.

Duplicative environmental reviews and inconsistencies in regulatory requirements among agencies make permitting of vegetation management projects difficult, increase costs, and slow the rate of progress of watershed restoration efforts. For example, Waste Discharge Requirement stipulations and conditions for Waivers of Waste Discharge Requirements vary among the nine RWQCBs. In some situations, projects require more than one permit related to water quality, sometimes from as many as three different agencies. A streamlined “one-stop shopping” approach would expedite projects and lower implementation costs.

Prescribed fires, which are being more widely used for vegetation management, are regulated by the California Air Resources Board (ARB) and local air pollution control agencies and can only be conducted on days approved for burning on the basis of air quality conditions. The USDA Forest Service is currently working cooperatively with the RWQCBs to increase opportunities for prescribed burning. Additionally, the ARB, along with the U.S. Environmental Protection Agency (EPA) and local air quality management districts, regulate biomass power plants that often utilize woody material generated by vegetation management projects.

Overarching Recommendations

The following recommendations are intended to address the issues identified in the previous section.

Monitoring and Research

Long-term monitoring is needed to understand hydrologic changes resulting from climate change and management actions, and more data collection stations are needed to accurately determine how changes in hydrology and water quality are related to climate change and forest management activities:

1. Additional stream gauges are needed throughout the forested regions of California to adequately represent the existing range of hydroclimatic and geologic conditions. In particular, gauges would be helpful on small (first to third order) reaches on unregulated and undiverted streams, in both managed and pristine watersheds.
2. Additional precipitation stations and snow courses are needed to increase the accuracy of determinations of climatic trends and evaluations of effects of management activities.
3. Additional water quality and sediment monitoring stations are needed to quantify the effects of climate change and forest management activities on surface water quality.
4. Additional long-term monitoring wells would be useful for understanding groundwater resources in forested watersheds.

Forest management for water resources could benefit from additional research on:

5. Effectiveness of BMPs in protecting beneficial uses of water.
6. Effects of vegetation and fuels management on soil moisture, groundwater recharge, and streamflow. More quantification of both the short- and long-term effects of prescribed fire on soil and water nutrient status is needed to determine the most beneficial and most ecosystem friendly return interval as a management strategy. Determination of the impacts of burn frequency on soil and vegetative properties that influence infiltration, percolation, surface runoff, and groundwater discharge would also be advantageous (Tahoe Science Consortium 2007).
7. Effects of wildfires and wildfire control measures on water quantity, water quality, and aquatic organisms.
8. Role and magnitude of groundwater storage in mountain meadows and its effects on streamflow regulation, and of the potential benefits of meadow restoration for water quantity and quality.
9. Sediment sources and erosion processes in managed and unmanaged forested watersheds.
10. Effects of riparian forests in maintaining stream temperatures and cycling nutrients.
11. Effects of urban trees in reducing non-point source pollution.

Coordination

Actions that would provide for better multi-party coordination of forest management, including communication between downstream water users and upstream forest managers, residents and workers, include:

12. Involvement of forest managers in integrated resource water management plan development.
13. Determination of mutually agreeable objectives for forest and meadow protection and restoration in terms of land area and timelines, and commitments from forest managers to meet these objectives.

14. Expanded authority and interagency agreements to allow federal, state, and non- governmental agencies to share expertise, staff time, and funding across jurisdictional boundaries for the purposes of watershed and water quality protection and improvement.
15. Develop a public education campaign directed at water users and communities in the Central Valley, Bay Area, and Southern California to increase support of forest management funding for improvement of water resources, particularly related to vegetation management.
16. Resolve water rights issues related to restoration of forested watersheds, and develop mechanisms for marketing of additional water made available by restoration projects.
17. Expand the scope of state water resource development and conservation measures to include headwaters areas of the state and urban forestry in metropolitan areas.
18. Increase eligibility of federal agencies for grant programs, and allow federal funds and in-kind services to be used as grant matches.

Regulatory Requirements

The water quality management plans developed by the SWRCB and forest management agencies can be revised to address concerns with impaired water bodies, while at the same time providing consistency and cost-effectiveness. Regulatory workloads can be reduced by combining environmental compliance into fewer streamlined procedures that would apply to all projects that meet criteria for low risk of adverse watershed effects or net beneficial water quality effects.

The following recommendations are directed at regulatory oversight of forest water resources:

19. Revise forest management agency water quality programs as necessary to identify, prioritize, and repair existing pollution sources, improve BMPs, and modify monitoring programs.
20. Incorporate existing Management Agency Agreements between the SWRCB and forest management agencies into cost-effective and consistent regulatory mechanisms compliant with current state law.
21. Deregulate low-risk noncommercial vegetation and fuels management projects that reduce the risks of catastrophic wildfires and therefore have net beneficial effects on water quality.
22. Complete a water quality management plan for the U.S. Bureau of Land Management.
23. Change the SWRCB's Water Quality Control Policy for Addressing Impaired Waters to incorporate Category 4B of EPA's Integrated Reporting Guidance, thereby allowing water quality management programs of other entities to be used to attain water quality standards in 303(d)-listed impaired waters in lieu of adopting total maximum daily loads (TMDLs) and duplicative TMDL implementation plans.

Beneficial Forest Management in Areas with Commercial-Scale Marijuana Cultivation

24. A combination of innovative prevention and enforcement approaches are needed to gain control over commercial-scale marijuana operations in California. Google Earth imagery and other remote sensing tools, such as infrared heat imaging, are now available to allow for increased site detection and information gathering. Even with these tools, however, new law enforcement strategies, a commitment to long-term investment of adequate resources, and large-scale changes in public policy are needed to change the current situation (Mallery 2011).
25. Education is also an important component of preserving public lands for public benefit, such as the production of abundant, clean water. Major knowledge gaps exist between the public,

politicians, and law enforcement agency personnel (Mallery 2011), although several recent newspaper stories, blog postings, and PowerPoint presentations to regulatory agencies and others have heightened general awareness of this threat to water quality in California (Dawson 2011). The effects of marijuana cultivation on water quality and fisheries resources were part of a state legislative hearing in Sacramento in February 2012, with discussion of possible legislative action. Water quality and fisheries protection are two essential components of a successful California Water Plan.

26. Commercial-scale marijuana cultivation on public and private lands is producing significant environmental problems. There are possible solutions, but without essential changes in law enforcement strategies and public policy, it is a problem that can be expected to continue into the foreseeable future (Mallery 2011).

Special Author's Note

This resource management strategy report is dedicated to Melvin Carmen of the North Fork Mono Tribe in 2009, in recognition of his vision of incorporating forest management into the California Water Plan.

References

References Cited

- Allen-Diaz BH. 1991. "Water table and plant species relationships in Sierra Nevada meadows." [Journal.] *American Midland Naturalist*. 126:30-43.
- Amaranthus M, Jubas H, Arthur D. 1989. "Stream shading, summer streamflow and maximum water temperature following intense wildfire in headwater streams." In: N. Berg N, editor. *Proceedings of the Symposium on Fire and Watershed Management, 26-28 October 1988*. Berkeley (CA): USDA Forest Service. Pacific Southwest Forest and Range Experiment Station. General Technical Report PSW-109. pp. 75-78.
- Backer DM, Jensen SE, McPherson GR. 2004. "Impacts of fire-suppression activities on natural communities." [Journal.] *Conservation Biology* 18(4): 937-946.
- Bales RC, Battles JJ, Chen Y, Conklin MH, Holst E, O'Hara K, Saksa P, Stewart W. 2011. *Forests and Water in the Sierra Nevada: Sierra Nevada Watershed Ecosystem Enhancement Project*. Report number 11.1. Merced (CA): Sierra Nevada Research Institute University of California, Merced. 39 pp.
- Barbour M, Pavlik BS, Lindstrom S. 1993. *California's Changing Landscapes: Diversity and Conservation of California Vegetation*. Sacramento (CA): The California Native Plant Society. 244 pp.
- Benavides-Solorio J, MacDonald LH. 2001. "Post-fire runoff and erosion from simulated rainfall on small plots, Colorado Front Range." [Journal.] *Hydrological Processes*. 15(15):2931-2952.

- Berlow EL, D'Antonio CM, Reynolds SA. 2002. "Shrub expansion in montane meadows: the interaction of local-scale disturbance and site aridity." [Journal.] *Ecological Applications*. 12(4):1103-1118.
- Beschta, RL 1990. "Effects of fire on water quantity and quality." In: Walstad JD, Radosevich SR, Sandberg DV editors. *Natural and Prescribed Fire in Pacific Northwest Forests*. Corvallis (OR): Oregon State University Press. pp. 219–232.
- Best DW, Kelsey HM, Hagans DK, Alpert M. 2004. "Role of fluvial hillslope erosion and road construction in the sediment budget of Garret Creek, Humboldt County, California." In: Nolan KM, Kelsey HM, Marron DC, editors. *Geomorphic Processes and Aquatic Habitat in the Redwood Creek Basin, Northwestern California*. Washington (DC): U.S. Geological Survey. Professional Paper 1454. pp. M1-M11
- Beyers JL 2004. "Postfire seeding for erosion control: Effectiveness and impacts on native plant communities." [Journal.] *Conservation-Biology* 18: 947-956.
- Blanchard FB. 1962. *Logging and Water Supply in the Mokelumne River Basin—A Preliminary Appraisal*. File Report. Oakland (CA): East Bay Municipal Utilities District, Water Resources and Planning Division. 3 pp.
- Brandow CA, Cafferata PH, Munn JR. 2006. *Modified Completion Report Monitoring Program: Monitoring Results from 2001 through 2004. Monitoring Study Group Final Report*. Sacramento (CA): California State Board of Forestry and Fire Protection. 80 pp.
- Cafferata PH, Munn JR. 2002. *Hillslope Monitoring Program: Monitoring Results from 1996 through 2001. Monitoring Study Group Final Report*. Sacramento (CA): California State Board of Forestry and Fire Protection. 114 pp.
- Cafferata PH 2005. *Herbert Avenue Meadow Restoration Project Hydrologic Review*. Memorandum dated May 25, 2005, sent to Mary Huggins, South Lake Tahoe, CA. California Department of Forestry and Fire Protection, Sacramento, CA. 12 p.
- Cafferata PH, Berbach M, Burke J, Hendrix J, Klamt R, Macedo R, Spittler T, Vyverberg K, Wright-Shacklett C. 2005. *Flood Prone Area Considerations in the Coast Redwood Zone. Final Report of the Riparian Protection Committee*. Sacramento (CA): California Department of Forestry and Fire Protection. 67 pp.
- Cafferata PH, Hall DO, Gentry GD. 2007a. "Applying scientific findings to forest practice regulations in California." In: *Proceedings of the 1997 NCASI West Coast Regional Meeting*. Special Report No. 97-13. 1997 NCASI West Coast Regional Meeting. 2007. Sept. 26-27, 1997. Portland (OR): National Council for Air and Stream Improvement. Research Triangle Park (NC): p. H-39 to H-46.
- Cafferata PH, Coe DBR, Harris RR. 2007b. "Water resource issues and solutions for forest roads in California." [Journal.] *Hydrological Science and Technology*. 23(1-4):39-56. Calder I, Hofer T,

- Vermont S, Warren P. 2007. "Towards a new understanding of forests and water." [Journal.] *Unasylva*. 229. 58:3-43.
- California Department of Forestry and Fire Protection. 2003a. *The Changing California: Forest and Range 2003 Assessment Technical Report, Forest Land Base*. Sacramento (CA): California Department of Forestry and Fire Protection, Fire and Resource Assessment Program. Chapter 2. Productive Capacity. Viewed online at: http://www.frap.cdf.ca.gov/assessment2003/Chapter2_Area/forestlandbase.html. Accessed: Nov. 10, 2009.
- . 2003b. *The Changing California: Forest and Range Assessment: Assessment Summary, Fire and Resource Assessment Program*. Sacramento (CA): California Department of Forestry Fire Protection. Variously paged. Viewed online at: <http://www.frap.cdf.ca.gov/assessment2003/index.html>.
- . 2010a. *California's Forests and Rangelands: 2010 Assessment*. Sacramento (CA): California Department of Forestry Fire Protection. Fire and Resource Assessment Program. 343 pp. Viewed online at: <http://frap.fire.ca.gov/assessment2010.html>.
- . 2010b. *California's Forests and Rangelands: 2010 Strategy Report*. Sacramento (CA): California Department of Forestry Fire Protection. Fire and Resource Assessment Program. 208 pp. Viewed online at: http://www.sierrainstitute.us/burneycreek-hatcreek/California_Forest_Strategy_Report_2010.pdf.
- . 2013. "Demonstration State Forests." Sacramento (CA): California Department of Forestry Fire Protection. [Web site.] Viewed online at: http://calfire.ca.gov/resource_mgt/resource_mgt_stateforests.php.
- California Department of Water Resources. 2008. *Managing an Uncertain Future: Climate Change Adaptation Strategies for California's Water*. Sacramento (CA): California Department of Water Resources.
- Campbell CG, Allen-Diaz B. 1997. "Livestock grazing and riparian habitat water quality: an examination of oak woodland springs in the Sierra foothills of California." In: Pillsbury NH, Verner, J, Tietje WD, editors. *Proceedings of a Symposium on Oak Woodlands: Ecology, Management, and Urban Interface Issues*. General Technical Report PSW-GTR-160. Mar. 19-22, 1996. San Luis Obispo (CA): USDA Forest Service. Pacific Southwest Research Station. Albany (CA): pp. 339-346. Viewed online at: <http://www.treesearch.fs.fed.us/pubs/28193>.
- Carroll EM, Miller WW, Johnson DW, Saito LS, Qualls RG, Walker RF. 2007. "Spatial analysis of a high magnitude erosion event following a Sierran wildfire." [Journal.] *Journal of Environmental Quality*. 36:1105-1111.
- Christensen GA, Campbell SJ, Fried JS, technical editors. 2008. *California's Forest Resources 2001–2005: Five-year Forest Inventory and Analysis Report*. General Technical Report PNW-GTR-763. Portland (OR): USDA Forest Service. Pacific Northwest Research Station. 183 pp.

- Coe DBR. 2006. *Sediment Production and Delivery from Forest Roads in the Sierra Nevada*. [Master's thesis.] Fort Collins (CO): Colorado State University. 110 pp.
- Cohen WL. 1982. *Effects of the Atlas Peak Fire on Surface Water Quality in Upper Milliken Creek Watershed, Napa County, California*. Master of Science Thesis. Davis (CA): University of California, Davis. Davis, CA. 132 pp.
- Conard SG, Beyers JL, Wohlgemuth PM. 1995. "Impacts of post-fire grass seeding on chaparral ecosystems: what do we know and where do we go from here?" Fairfield (WA): In: Keeley JE, Scott T editors. *Brushfires in California Wildlands: Ecology and Resource Management*. International Association of Wildland Fire. pp. 149-161.
- Cook C, Dresser A. 2007. "Erosion and channel adjustments following forest road decommissioning Six Rivers National Forest." In: Furniss M, Clifton C, Ronnenberg K, editors. *Advancing the Fundamental Sciences: Proceeding of the Forest Service National Earth Sciences Conference*. Forest Service National Earth Sciences Conference. Oct. 18-2, 2004. San Diego (CA): USDA Forest Service. Pacific Northwest Research Station. Portland (OR): pp. 297-309.
- Cornwell K, Brown K. 2008. *Physical and Hydrological Characterization of Clark's Meadow in the Last Chance Watershed of Plumas County*. Sacramento (CA): Report to the Natural Heritage Institute Mountain Meadows IRWMP. California State University Sacramento. Department of Geology. 38 pp.
- Cram DS, Baker TT, Fernald AG, Madrid A, Rummer B. 2007. "Mechanical thinning impacts on runoff infiltration and sediment yield following fuel reduction treatments in a southwestern dry mixed conifer forest." [Journal.] *Journal of Soil and Water Conservation*. 62(5):359-366.
- California State Board of Forestry and Fire Protection. 1996. "Assets at risk and their role in the fire plan: water and watersheds." In: *California Fire Plan: A Framework for Minimizing Costs and Losses from Wildland Fires*. Sacramento (CA): California State Board of Forestry and Fire Protection. pp. 73-82.
- Council of Western State Foresters. 2007. *Forestry Best Management Practices for Western States: A Summary of Approaches to Water Quality Implementation and Effectiveness Monitoring. Final Report*. Lakewood (CO): Council of Western State Foresters. 20 pp. Viewed online at: http://www.wflcenter.org/news_pdf/240_pdf.pdf.
- Dailey S, Fites JA, Reiner A, Mori S. 2008. *Fire Behavior and Effects in Fuel Treatments and Protected Habitat on the Moonlight Fire*. Vallejo (CA): USDA Forest Service, Pacific Southwest Region. Adaptive Management Service Enterprise Team. 60 pp. Viewed online at: http://www.fs.fed.us/r5/hfqlg/monitoring/resource_reports/fire_and_smoke/dfpz_effectiveness/moonlight_fire_effects_assessment.pdf. Accessed Nov. 10, 2009.
- Dawson B. 2011. "Pollute, poach, dewater: legacies of illicit cannabis grows." Green Blog - UC work in natural resources, pest management, climate change and sustainable agriculture. Oakland (CA):

- University of California Agriculture and Natural Resources. [Blog.] Nov., 16. Viewed online at: http://ucanr.org/sites/UCCNP/Green_Blog/?blogpost=6188&blogasset=18704.
- Dombeck M. 2007. "Deterioration of forest roads threatens our drinking water." San Jose Mercury News. San Jose (CA): San Jose Mercury News. [Web site.] Viewed online at: http://www.mercurynews.com/opinion/ci_7211103. Accessed Jan. 2012.
- Driscoll DG, Carter JM, Ohlen DO. 2004. *Hydrologic Effects of the 1988 Galena Fire, Black Hills Area, South Dakota*. Denver (CO): Water-Resources Investigations Report 03-4323. U.S. Geological Survey. 67 pp. Viewed online at: <http://pubs.water.usgs.gov/wri034323>.
- Dudley N, Stolton S. 2003. *Running Pure: The Importance of Forest Protected Areas to Drinking Water*. Washington (DC): World Bank/WWF Alliance for Forest Conservation and Sustainable Use. 114 pp. Viewed online at: http://www.panda.org/what_we_do/how_we_work/conservation/forests/publications/?uNewsID=8443.
- Dwire KA, Kauffman JB, Baham JE. 2006. "Plant species distribution in relation to water-table depth and soil redox potential in montane riparian meadows." [Journal.] *Wetlands*. 26(1):131-146.
- Elliot WJ. 2010. "Effects of forest biomass use on watershed processes in the western United States." [Journal.] *Western Journal of Applied Forestry*. 25(1):12-17.
- Fiori RA. 2005. *Instream and Riparian Dynamics Following Wildfire in an Alluvial Old-growth Redwood Forest--Canoe Creek, Humboldt Redwoods State Park*. California State Parks. North Coast Redwoods District. Eureka (CA): 22 pp.
- Fryoff-Hung A, Viers JH. 2013. *Sierra Nevada Meadow Hydrology Assessment, Final Project Report*. University of California, Davis, 1114 pp. Viewed online at <http://meadows.ucdavis.edu/data/9>.
- Furniss MJ, Roelofs TD, Yee CS. 1991. "Road construction and maintenance." In: Meehan WR, editor. *Influences of Forest and Rangeland Management*. Bethesda (MD): Special Publication 19. American Fisheries Society. pp. 297-324.
- Furniss MJ, Ledwith TS, Love MA, McFadin BC, Flanagan SA. 1998. *Response of Road-Stream Crossings to Large Flood Events in Washington, Oregon and Northern California*. Washington (DC): USDA Forest Service. Technology and Development Program. 14 pp.
- Giusti GA. 2012. Illicit cannabis production impacts on forest and wildland resources. Sacramento (CA): Presentation to the California State Board of Forestry and Fire Protection. Apr. 4, 2012. [PowerPoint presentation.] Viewed online (2010 version of the presentation) at: <http://ucanr.org/blogs/Green/blogfiles/9273.pdf>.
- Gokbulak F, Serengil Y, Ozhan S, Ozyuvaci N, Balç, N. 2008. "Effect of timber harvest on physical water quality characteristics." [Journal.] *Water Resources Management*. 22(5):635-649.

- Grace JM, Skaggs RW, Chescheir GM. 2006. "Hydrologic and water quality effects of thinning loblolly pine." [Journal.] *Transactions of the American Society of Agricultural Engineers*. 49(3):645-654.
- Hammersmark CT, Rains MC, Mount JF. 2008. "Quantifying the hydrological effects of stream restoration in a montane meadow, Northern California USA." *River Research and Applications* Vol. 24 (6): 735-753. [Journal]. Viewed online at: <http://onlinelibrary.wiley.com/doi/10.1002/rra.1077/abstract>.
- Hammersmark CT, Dobrowski SZ, Rains MC, Mount JF. 2010. "Simulated Effects of Stream Restoration on the Distribution of Wet-Meadow Vegetation." [Journal.] *Restoration Ecology*. 18:882–893.
- Harris RR. 2011. *Policies and Procedures for Protecting Anadromous Salmonid Habitat in Del Norte, Humboldt, Mendocino, Trinity and Siskiyou Counties*. Final Report prepared for the Five Counties Salmonid Conservation Program. Weaverville (CA): 52 pp. Viewed online at: http://www.5counties.org/docs/2011_reassessment.pdf.
- Harris RR, Cafferata PH. 2005. *Forest Loss and Fragmentation; Session 2: Impacts and Implications—Impacts on water quantity and quality* [Abstract]. California Forest Futures 2005 Conference, May 24-25, 2005, Sacramento, CA.
- Hatchett B, Hogan MP, Grismer ME. 2006. "Mechanical mastication thins Lake Tahoe forest with few adverse impacts." [Journal.] *California Agriculture*. 60(2):77-82.
- Heard AM. 2005. *Effects of Landscape Scale Prescribed Fire on Hydrology and Stream Chemistry*. Master of Science Thesis. Fort Collins (CO): Colorado State University. 119 pp.
- Hill BR. 1990. "Groundwater discharge to a headwater valley northwestern Nevada USA." [Journal.] *Journal of Hydrology*. 113:265-283.
- Hill BR, Mitchell-Bruker S. 2010, Comment on "A framework for understanding the hydroecology of impacted wet meadows in the Sierra Nevada and Cascade Ranges, California, USA." [Comment on paper published in *Hydrogeology Journal* 17:229–246. 2009, by Loheide II SP, Deitchman RS, Cooper DJ, Wolf EC, T. Hammersmark CT, Lundquist JD.] *Hydrogeology Journal* 18(7):1741-1743.
- Hoffman J, Roby K, Bohm B. 2013, *Effects of Meadow Restoration on Stream Flow in the Feather River Watershed: A Review Based on Monitoring Data and Pertinent Research, Version 5.5 (final)*, June 12, 2013, report to Feather River Coordinated Resource Management Council Executive Committee, 33 pp.
- Hubbert K, Busse M, Overby S, Shestak C. 2010. Soil heating beneath burn piles: effects on soil physical properties and water chemistry in the Lake Tahoe Basin. [Poster presentation.] ASA, CSSA, SSSA 2010 International Annual Meeting, Oct. 21-Nov 4, 2010. Long Beach (CA): USDA Forest Service. Pacific Southwest Research Station.

- Hughes JE. 1934. *Erosion Control Progress Report*. Plumas National Forest. Milford Ranger District. 26 pp.
- Ice G, Dent L, Robben J, Cafferata P, Light J, Sugden B, Cundy T. 2004. "Programs Assessing Implementation and Effectiveness of State Forest Practice Rules and BMPs in the West." *Water, Air, and Soil Pollution: Focus* 4(1): 143-169.
- Ice G, Shilling EG, Vowell JG. 2010. "Trends for Forestry Best Management Practices Implementation." *Journal of Forestry*. 108. 267-273.
- Jewett DG, Lord ML, Miller JR, Chambers JC. 2004. "Geomorphic and hydrologic controls on surface and subsurface flow regimes in riparian meadow ecosystems." In: Chambers JC, Miller JR, editors. *Great Basin Riparian Ecosystems*. Covelo (CA): Society for Ecological Restoration International. Island Press. Chapter 5, pp. 124-161.
- Jones JA, Grant GE. 1996. "Peak flow responses to clear-cutting and roads in small and large basins, western Cascades, Oregon." [Journal.] *Water Resources Research*. 32(4):959-974.
- Kattelman R, Embury M. 1996. "Riparian areas and wetlands." In: *Sierra Nevada Ecosystem Project. Final Report to Congress, Volume III: Assessments, Commissioned Reports and Background Information*. Davis (CA): University of California, Centers for Water and Wildland Resources. Chapter 5, pp. 201-274. Viewed online at: http://pubs.usgs.gov/dds/dds-43/VOL_III/VIII_TOC.PDF.
- Kellar G, Sherar J. 2003. *Low Volume Roads Engineering: Best Management Practices Field Guide*. Washington (DC): Prepared for U.S. Agency for International Development by the Conservation Management Institute, Virginia Polytechnic Institute and State University and USDA Forest Service. Viewed online at: http://pdf.usaid.gov/pdf_docs/PNADB595.pdf.
- Korte A, MacDonald LH. 2007. *Preliminary Summary: Sediment Production and Delivery in the Southern Sierra Nevada*. Watershed Science Program, Colorado State University, Fort Collins (CO). 5 pp. [Unpublished paper.]
- Kraebel CJ, Pillsbury AF. 1934, *Handbook of Erosion Control in Mountain Meadows*. USDA Forest Service. California Forest and Range Experiment Station. Berkeley(CA): 69 pp.
- Krieger DJ. 2001. *The Economic Value of Forest Ecosystem Services: A Review*. Washington (DC): The Wilderness Society. 40 pp. Viewed online at: http://www.cfr.washington.edu/classes/esrm.465/2007/readings/WS_valuation.pdf.
- Lautz LK. 2008. "Estimating groundwater evapotranspiration rates using diurnal water-table fluctuations in a semi-arid riparian zone." [Journal.] *Hydrogeology Journal*. 16: 483-497.
- Liang L, Kavvas ML, Chen ZQ, Anderson M, Ohara N, Wilcox J, Mink L. 2007. "Modeling river restoration impact on flow and sediment in a California watershed." In: Kabbes KC, editor. *Proceedings of the 2007 World Environmental and Water Resources Congress*. 2007 World

- Environmental and Water Resources Congress. May 15-19, 2007. Tampa (FL): American Society of Civil Engineers. Reston (VA): pp. 1-10. Viewed online at: [http://ascelibrary.org/doi/abs/10.1061/40927\(243\)353](http://ascelibrary.org/doi/abs/10.1061/40927(243)353).
- Ligon F, Rich A, Ryneerson G, Thornburgh D, Trush, W. 1999. *Report of the Scientific Review Panel on California Forest Practice Rules and Salmonid Habitat*. Sacramento (CA): California Natural Resources Agency and the National Marine Fisheries Service. 181 pp.
- Loheide SPII, Gorelick SM. 2005. "A local-scale, high-resolution evapotranspiration mapping algorithm (ETMA) with hydroecological applications at riparian meadow restoration sites." [Journal.] *Remote Sensing of Environment*. 98: 182-200.
- Loheide SP II, Gorelick SM. 2007. "Riparian hydroecology: a coupled model of the observed interactions between groundwater flow and meadow vegetation patterning." [Journal.] *Water Resources Research*. 43:W07414. Viewed online at: <http://onlinelibrary.wiley.com/doi/10.1029/2006WR005233/abstract>.
- Liquori M, Cafferata P, Boston K, Gienger R, Hope D. 2012. "The VTAC Committee: developing guidance for an alternative regulatory pathway to the anadromous salmonid protection rules." In: Standiford RB, Weller TJ, Piirto DD, Stuart, JD technical coordinators. *Proceedings of Coast Redwood Forests in a Changing California: A Symposium for Scientists and Managers*. Albany (CA): USDA U.S. Forest Service. Pacific Southwest Research Station General Technical Report PSW-GTR-238. pp 79-89.
- McCashion JD, Rice RM. 1983. "Erosion on logging roads in northwestern California: How much is avoidable?" [Journal.] *Journal of Forestry*. 81(1):23-26.
- McClurkin DC, Duffy PD, Nelson NS. 1987. "Changes in forest floor and water quality following thinning and clearcutting of 20-year-old pine." [Journal.] *Journal of Environmental Quality*. 16:237-241.
- MacDonald LH, Coe D, Litschert S. 2004. "Assessing cumulative watershed effects in the central Sierra Nevada: hillslope measurements and catchment-scale modeling." In: Murphy DD, Stine PA, editors. *Proceedings of the Sierra Nevada Science Symposium*. Sierra Nevada Science Symposium. Oct. 7-10, 2002, Kings Beach (CA): USDA Forest Service. Pacific Southwest Research Station. Albany (CA): pp. 149-157. Viewed online at: <http://www.treesearch.fs.fed.us/pubs/26449>.
- MacDonald LH, Brown E, Pietraszek JH, Welsh M, Eccleston D, Larsen IJ. 2006. "Long-term Sediment Production and Recovery Rates from Forest Thinning, Roads, and Wildfires." American Geophysical Union, Fall Meeting 2006, Abstract #B22E-08.
- MacDonald LH, Coe, D. 2007. "Influence of headwater streams on downstream reaches in forested areas." [Journal.] *Forest Science*. 53(2):148-168.

- Maco S, McPherson G, Simpson J, Peper P, Xiao Q. 2003. *City of San Francisco, California Street Tree Resource Analysis*. Center for Urban Forest Research, Pacific Southwest Research Station, USDA Forest Service. CUF3-3.
- McPherson G, Simpson J, Peper P, Maco S, Xiao Q. 2005. "Municipal forest benefits and costs in five U.S. cities." [Journal.] *Journal of Forestry*. (Dec):411-416.
- Madej MA. 2001. "Erosion and sediment delivery following removal of forest roads." [Journal.] *Earth Surface Processes and Landforms*. 26:1-16.
- Madrid A, Fernald AG, Baker TT, Vanleeuwen DM. 2006. "Evaluation of silvicultural treatment effects on infiltration, runoff, sediment yield, and soil moisture in a mixed conifer New Mexico forest." [Journal.] *Journal of Soil and Water Conservation*. 61(3):159-168.
- Mahlum SK, Eby LA, Young MK, Clancy CG, Jakober M. 2011. "Effects of wildfire on stream temperatures in the Bitterroot River Basin, Montana." [Journal.] *International Journal of Wildland Fire*. 20:240-247.
- Mallery M. 2011. "Marijuana national forest: encroachment on California public lands for cannabis cultivation." Office of Undergraduate Research, University of California, Berkeley. Berkeley Undergraduate Journal. 23(2). 50 pp. Viewed online at: <http://escholarship.org/uc/item/7r10t66s>.
- Martin S. 2006. Topics in mountain meadows. Merced (CA): [PowerPoint class presentation.] University of California, Merced. 45 pp.
- Martinson EJ, Omi PN. 2003. "Performance of fuel treatments subjected to wildfires." In: Omi PN, Joyce LA, editors. *Fire, Fuel Treatments, and Ecological Restoration: Conference Proceedings, April 16-18, 2002*. Fort Collins (CO): Proceedings RMRS-P-29. USDA Forest Service. Rocky Mountain Research Station. pp. 7-13
- Meixner T, Wohlgemuth P. 2004. "Wildfire impacts on water quality." [Journal.] *Southwest Hydrology* 3(5):24-25.
- Micheli ER, Kirchner JW. 2002. "Effects of wet meadow riparian vegetation on streambank erosion. 2. Measurements of vegetated bank strength and consequences for failure mechanics." [Journal.] *Earth Surface Processes and Landforms*. 27:687-697.
- Miller WW, Johnson DW, Loupe TM, Sedinger JS, Carroll EM, Murphy JD, Walker RF, Glass D. 2006. "Nutrients flow from runoff at burned forest site in Lake Tahoe Basin." [Magazine article] *California Agriculture*. University of California Agriculture and Natural Resources. Apr-Jun:65-71.
- Moody JA, Martin DA. 2001. "Post-fire, rainfall intensity-peak discharge relations for three mountainous watersheds in the western USA. [Journal.] *Hydrological Processes* 15: 2981-2993.

- Moody JA, Martin DA, Cannon SH. 2008. "Post-wildfire erosion response in two geologic terrains in the western USA." [Journal.] *Geomorphology*. 95:103-118.
- Moore RD, Wondzell SM. 2005. "Physical hydrology and the effects of forest harvesting in the Pacific Northwest: a review." [Journal.] *Journal of the American Water Resources Association*. Aug.:763-784.
- Moser E. 2007. *Aerial Application of Fire Retardant Environmental Assessment: Hydrology Specialist Report*. USDA Forest Service. T.E.A.M.S. Enterprise Unit. 10 pp. Viewed online at: <http://www.fs.fed.us/fire/retardant/references/01Hydro.pdf>.
- Murphy K, Rich T, Sexton T. 2007. *An Assessment of Fuel Treatment Effects on Fire Behavior, Suppression Effectiveness, and Structure Ignition on the Angora Fire*. Vallejo (CA): USDA Forest Service. R5-TP-025. 32 pp.
- Naiman RJ, Fetherston KL, McKay SJ, Chen J. 1998. "Riparian forests." In: Naiman RJ, Bilby RE, editors. *River Ecology and Management: Lessons from the Pacific Coastal Ecoregion*. New York: Springer-Verlag. pp. 289-323. [Book.]
- National Academy of Sciences. 2008. *Hydrologic Effects of a Changing Forest Landscape*. Washington (DC): Report in Brief. Report brief was prepared by the National Research Council. Viewed online at: http://dels.nas.edu/resources/static-assets/materials-based-on-reports/reports-in-brief/forest_hydrology_final.pdf.
- National Interagency Fire Center (. 2010. "Suppression chemicals and delivery systems." In: Interagency Standards for Fire and Fire Aviation Operations ("Red Book"). Boise (ID): Viewed online at: <http://www.nifc.gov/PUBLICATIONS/redbook/2011/Ch12.pdf>. Chapter 12. pp 12-1—12-6.
- National Wildfire Coordinating Group Training Working Team. 2004. *Resource Advisor's Guide for Wildland Fire*. PMS 313. NFES 1831. Boise (ID): National Interagency Fire Center. 14 pp. Viewed online at: http://www.nwcg.gov/pms/pubs/RAGuide_2004.pdf.
- Neariy DG, Ryan KC, DeBano LF, editors. 2005. *Wildland Fire in Ecosystems: Effects of Fire on Soils and Water*. General Technical Report RMRS-GTR-42, Vol. 4. Ogden (UT): USDA Forest Service, Rocky Mountain Research Station. 250 pp.
- Norman S, Loupe T, Keely J. 2008. *Heavenly Creek SEZ Demonstration Project 2007 Soil Monitoring Report*. South Lake Tahoe (CA): USDA Forest Service. Lake Tahoe Basin Management Unit. 58 pp. Viewed online at: http://www.fs.usda.gov/Internet/FSE_DOCUMENTS/fsm9_045987.pdf.
- Novak DJ, Greenfield EJ. 2009. *Urban and Community Forests of the Pacific Region: California, Oregon, Washington*. GTR NRS-65. St. Paul (MN): USDA Forest Service, Northern Research Station. 43 pp.
- Ohara N, Kavvas, ML, Chen ZQ, Liang L, Anderson M, Wilcox J, Mink L. 2013. *Modelling Atmospheric and Hydrologic Processes for Assessment of Meadow Restoration Impact on Flow and Sediment*

- in a Sparsely Gauged California Watershed*. Hydrological Processes. DOI: 10.1002/hyp.9821.,14 pp.
- Omi PN, Martinson, EJ. 2004. "Effectiveness of thinning and prescribed fire in reducing wildfire severity." In: Murphy DD, Stine PA editors. *Proceedings of the Sierra Nevada Science Symposium: Science for Management and Conservation*. Albany (CA): General Technical Report PSW-193. USDA Forest Service. pp. 87-92
- Onda Y, Dietrich WE, Booker F. 2007. "Evolution of overland flow after a severe forest fire Point Reyes California." [Journal.] *Catena*. 72:13-20.
- Pacific Watershed Associates. 2005. *Evaluation of Road Decommissioning CDFG Fisheries Restoration Grant Program 1998 to 2003*. Arcata (CA): Pacific Watershed Associates. Final Report submitted to the California Department of Fish and Game under Contract #P0210559. 91 pp. Viewed online at: http://www.pacificwatershed.com/sites/default/files/CDFG_PWA_decom_report.pdf.
- Peterson DH, Stewart I, Murphy F. 2008. "Principle hydrologic responses to climatic and geologic variability in the Sierra Nevada, California." [Journal.] *San Francisco Estuary and Watershed Science*. California Bay-Delta Authority Science Program and John Muir Institute of the Environment. 6(1): Article 3.
- Potts DF, Peterson DL, Zuuring HR. 1989. *Estimating Postfire Water Production in the Pacific Northwest*. Berkeley (CA): USDA Forest Service. Research Paper PSW-197. USDA Forest Service. Pacific Southwest Forest and Range Experiment Station. 9 pp.
- Ratliff RD. 1985. *Meadows in the Sierra Nevada of California: State of Knowledge*. Berkeley (CA): General Technical Report PSW-84. USDA Forest Service. Pacific Southwest Forest and Range Experiment Station. 52 pp. Viewed online at: <http://www.treesearch.fs.fed.us/pubs/27313>.
- Rector JR, MacDonald LH 1987. "Water yield opportunities on National Forest lands in the Pacific Southwest Region." In: *Proceedings, California Watershed Management Conference*. Report No. 11. Wildland Resources Center. University of California, Berkeley. pp. 68-73.
- Reid LM. 2012. "Comparing hydrologic responses to tractor-yarded selection and cable-yarded clearcut logging in a coast redwood forest." In: Standiford RB, Weller TJ, Piirto DD, Stuart JD technical coordinators. *Proceedings of the Coast Redwood Forests in a Changing California: A Symposium for Scientists and Managers*. Albany (CA): General Technical Report PSW-GTR-238. USDA Forest Service. Pacific Southwest Research Station. pp. 141- 151.
- Reneau SL, Katzman D, Kuyumjian GA, Lavine A, Malmon DV. 2007. "Sediment delivery after a wildfire." [Journal.] *Geology*. 35(2):151-154.
- Robichaud PR. 2000. "Forest fire effects on hillslope erosion: what we know." Portland (OR): Watershed Management Council (WMC) Archives 9(1). Winter 2000. at Watershed.org. Viewed online at: <http://www.watershed.org/?q=node/329>. Accessed: Nov. 12, 2009.

- Robichaud PR, Beyers JL, Neary DG. 2000. *Evaluating the Effectiveness of Postfire Rehabilitation Treatments*. Fort Collins (CO): General Technical Report RMRS-GTR-63. USDA Forest Service. Rocky Mountain Research Station. 89 pp. Viewed online at: http://www.fs.fed.us/rm/pubs/rmrs_gtr063.pdf.
- Robichaud PR, Pierson FB, Brown, RE. 2007. "Runoff and erosion effects after prescribed fire and wildfire on volcanic ash-cap soils." In: Page-Dumroese D. Miller R. Mital J. McDaniel P. Miller D, editors. *Volcanic-Ash-Derived Forest Soils of the Inland Northwest*. Fort Collins (CO): USDA Forest Service Proceedings RMRS-P-44. Properties and Implications for Management and Restoration. Nov. 9-10, 2005. Coeur d'Alene (ID): USDA Forest Service. Rocky Mountain Research Station. pp. 83-94. Viewed online at: <http://www.treesearch.fs.fed.us/pubs/26215>.
- Robichaud PR, MacDonald LH, Foltz, RB. 2010. "Fuel management and erosion." In: Elliot WJ, Miller IS, Audin L., editors. *Cumulative Watershed Effects of Fuel Management in the Western United States*. Fort Collins: (CO): USDA, Forest Service, Rocky Mountain Research Station. Chapter 5 pp. 79-100. General Technical Report RMRS-GTR-231.
- Rowe PB, Countryman CM, Storey HC. 1949. *Probable Peak Discharges and Erosion Rates from Southern California Watersheds as Influenced by fire*. Berkeley (CA): USDA Forest Service. California Forest and Range Experiment Station 342 pp.
- . 1954. *Hydrologic Analysis Used to Determine Effects of Fire on Peak Discharge and Erosion Rates in Southern California Watersheds*. Berkeley (CA): USDA. California Forest and Range Experiment Station.
- Rulli MC, Rosso R. 2007. "Hydrologic response of upland catchments to wildfires." [Journal.] *Advances in Water Resources*. 30(10):2072-2086.
- Stephens SL, Meixner T, Poth M, McGurk B, Payne D. 2004. "Prescribed fire, soils, and stream water chemistry in a watershed in the Lake Tahoe Basin." [Journal.] *International Journal of Wildland Fire* 13:27-35.
- Stephens SL, Moghaddas JJ. 2005. "Experimental fuel treatment impacts on forest structure potential fire behavior and predicted tree mortality in a California mixed conifer forest." [Journal.] *Forest Ecology and Management*. 215:21- 36.
- Stephens SL, Martin RE, Clinton NE. 2007. "Prehistoric fire area and emissions from California's forests, woodlands, shrublands and grasslands." [Journal.] *Forest Ecology and Management*. 251:205-216.
- Stohlgren TJ, DeBenedetti SH, Parsons DJ. 1989. "Effects of herbage removal on productivity of selected high-Sierra meadow community types." [Journal.] *Environmental Management*. 13(4):485-491.
- Tague C, Heyn K, Christiansen L. 2009. "Topographic controls on spatial patterns of conifer transpiration and net primary productivity under climate warming in mountain ecosystems." [Journal.] *Ecohydrology* 2: 541-554.

- Tague C, Valentine S, Kotchen M. 2008. "Effect of geomorphic channel restoration on streamflow and groundwater in a snowmelt-dominated watershed." [Journal.] *Water Resources Research* 44(W10415): 10 pp.
- Tahoe Science Consortium. 2007. *Comprehensive Science Plan for the Lake Tahoe Basin: Conceptual Framework and Research Strategies*. Final draft prepared as a deliverable under U.S. Environmental Protection Agency Cooperative Agreement X7-96963901-0. 290 pp;
- Thorsen K. 2011. "Exploring the problem of domestic marijuana cultivation." U.S. Senate Caucus on International Narcotics Control. Statement for the Record by Kim Thorsen, Deputy Assistant Secretary, Law Enforcement, Security and Emergency Management. Department of the Interior. Dec. 7. [Statement for the Record.] Viewed online at: <http://drugcaucus.senate.gov/hearing-12-7-11/DOI%20Statment%20for%20the%20Record%20on%20Marijuana%20Cultivation.pdf>.
- Tree City USA. 2010. "How trees can retain stormwater runoff." *Tree City USA Bulletin*. No. 55. pp. 1-8. Nebraska City (NE): Published for the Friends of Tree City USA by Arbor Day Foundation. [E-news.] Viewed online at: http://www.fs.fed.us/psw/programs/uesd/uep/products/11/800TreeCityUSABulletin_55.pdf.
- Troendle CA, Nankervis JM, Peavy A. 2007. *Impacts of Vegetation Management on Water Yield. Final Report*. Fort Collins (CO): The Herger-Feinstein Quincy Library Group Project. USDA Forest Service. Pacific Southwest Region. Resource Reports - Hydrology and Fisheries. Contract AG 3187 D 05 0043. 23 pp. Viewed online at: http://www.fs.fed.us/r5/hfqlg/monitoring/resource_reports/hydrology_and_fisheries/Water_Yield_Full_Report_Feb242007.pdf.
- Troendle CA, MacDonald LH, Luce CH, Larsen IJ. 2010. "Fuel management and water yield." In: Elliot WJ Miller IS, Audin L. editors. *Cumulative Watershed Effects of Fuel Management in the Western United States*. Fort Collins (CO): USDA. U.S. Forest Service Rocky Mountain Research Station. General Technical Report RMRS-GTR-231. Chapter 7. pp. 126-148.
- Turner KM. 1991. "Water salvage from Mediterranean-type ecosystems." In: Bailey HE, Forrest C, Snow L, editors. *Water Supply and Water Reuse: 1991 and Beyond. Proceedings of the American Water Resources Association Annual Symposium*. American Water Resources Association Annual Symposium. Jun. 2-6, 1991. San Diego (CA): American Water Resources Association. [Book.] pp. 83-90.
- USDA U.S. Forest Service. 1999. *Benefits of the Urban Forest*. Davis (CA): Fact Sheet #1. USDA Forest Service. Pacific Southwest Research Station. Center for Urban Forest Research. 2 pp.
- . 2000. *Water and the Forest Service*. Washington (DC): FS-660. USDA U.S. Forest Service. 40 pp. Viewed online at: <http://www.fs.fed.us/publications/policy-analysis/water.pdf>.
- . 2001. Sierra Nevada Framework Planning Amendment FEIS. USDA U.S. Forest Service. Vallejo (CA): [Web site.] Viewed online at: <http://www.fs.usda.gov/detail/r5/landmanagement/planning/?cid=stelprdb5415847>.

- . 2004. *Best Management Practices Evaluation Program: 1992-2002 Monitoring Results*. Vallejo (CA): USDA Forest Service. Pacific Southwest Region. Final Report. 76 pp.
- . 2005. *A Strategic Assessment of Forest Biomass and Fuel Reduction Treatments in Western states*. Fort Collins (CO): 17 p. General Technical Report RMRS-GTR-149. USDA Forest Service. Rocky Mountain Research Station. Viewed online at: http://www.fs.fed.us/rm/pubs/rmrs_gtr149.pdf.
- . 2006. *Urban Watershed Forestry Manual*. Viewed online at: www.na.fs.fed.us/watershed. Accessed: Aug., 2011.
- . 2007a. *An Assessment of Fuel Treatment Effects on Fire Behavior, Suppression Effectiveness, and Structure Ignition on the Angora Fire*. Vallejo (CA): R5-TP-025. USDA Forest Service. Pacific Southwest Research Station. Viewed online at: <http://www.cnpsd.org/fire/angorafireusfsfullreport.pdf>. Accessed: Nov. 10, 2009.
- . 2007b. *Land Areas of the National Forest System*. Washington (DC): FS-383. USDA Forest Service. 131 pp.
- . 2008. *Land Areas of the National Forest System as of September 30, 2007*. Washington (DC): FS-383. USDA Forest Service. 131 pp.
- . 2009. *Best Management Practices Evaluation Program: 2003-2007 Monitoring Results. Final Report*. Vallejo (CA): USDA Forest Service. Pacific Southwest Region. 28 pp.
- . 2011a. *Responding to Climate Change in National Forests: A Guidebook for Developing Adaptation Options*. Portland (OR): General Technical Report PNW-GTR-855. USDA Forest Service. Pacific Northwest Research Station. Viewed online at: <http://www.fs.fed.us/psw/publications/millar/Peterson%20et%20al.%20%28Adaptation%20Guidbook%29%20%28Feb.%202011%29.pdf>.
- . 2011b. *National Report on Sustainable Forests – 2010*. Washington (DC): FS-979. USDA Forest Service. Viewed online at: <http://www.fs.fed.us/research/sustain/national-report.php>.
- . 2011c. “Operation Full Court Press concludes three week operation targeting illegal marijuana plants on public lands.” USDA Forest Service Pacific Southwest Region press release dated August 8, 2011. Viewed online at: <http://www.fs.fed.us/r5/news/2011/OperationFullCourtPress.shtml>.
- Van de Water K, North M. 2011. “Stand structure, fuel loads, and fire behavior in riparian and upland forests, Sierra Nevada Mountains, USA; a comparison of current and reconstructed conditions.” [Journal.] *Forest Ecology and Management* 262: 215-228.
- Weaver WE, Hagans DK. 1994. *Handbook for Forest and Ranch Roads: A Guide for Planning, Designing, Constructing, Reconstructing, Maintaining, and Closing Wildland Roads*. Ukiah (CA): Mendocino County Resource Conservation District. 161 pp.

- Weaver W, Hagans D, Weppner E. 2006. "Upslope erosion inventory and sediment control guidance." Part X. In: *California Salmonid Stream Habitat Restoration Manual*. 1998 revised 3rd edition. Sacramento (CA): California Department of Fish and Game. 108 pp.
- Welsch HH. 2011. "Frogs, fish and forestry: an integrated watershed network paradigm conserves biodiversity and ecological services." [Journal.] *Diversity* 3: 503-530.
- Wilzbach MA, Harvey BC, White JL, Nakamoto RJ. 2005. "Effects of riparian canopy opening and salmon carcass addition on the abundance and growth of resident salmonids." [Journal.] *Canadian Journal of Fisheries and Aquatic Sciences*. 62: 58-67.
- Weixelman DA, Hill B, Cooper DJ, Berlow EL, Viers JH, Purdy SE, Merrill AG, Gross SE. 2001, *Meadow Hydrogeomorphic Types for the Sierra Nevada and Southern Cascade Ranges in California*. USDA Forest Service. R5-TP-034. 33 pp. Wohlgemuth PM, Beyers JL, Wakeman CD, Conard SG. 1998. "Effects of fire and grass seeding on soil erosion in southern California chaparral." In: Gray S Chairman. *Proceedings of the 19th Annual Forest Vegetation Management Conference: Wildfire Rehabilitation*. Redding (CA) University of California Cooperative Extension Forestry. pp. 41-51
- Wohlgemuth PM. 2001. "Prescribed fire as a sediment management tool in southern California chaparral watersheds." In: *Proceedings of the 7th Federal Interagency Sedimentation Conference*. March 25-29. Reno (NV): Federal Energy Regulatory Commission. 2:X-49-X-56.
- Wohlgemuth PM, Beyers JL, Hubbert KR. 2009. "Rehabilitation Strategies after Fire: The California, USA experience. pp. 511-536 In: Cerdá A, Robichaud PR editors. *Fire Effects on Soils and Restoration Strategies*. Enfield (NH): Science Publishers.
- Wondzell SM. 2001. "The influence of forest health and protection treatments on erosion and stream sedimentation in forested watersheds of eastern Oregon and Washington. [Journal.] *Northwest Science* 75:128-140.
- Wood SH. 1975. "Holocene stratigraphy and chronology of mountain meadows Sierra Nevada California." In: *Earth Surface Monograph*. Vallejo (CA): Volume 4. USDA Forest Service. Pacific Southwest Region. 180 pp.
- Xiao Q, McPherson G, Simpson J, Ustin S. 1998. "Rainfall interception by Sacramento's urban forest." [Journal.] *Journal of Arboriculture*. 24(4):235-244.
- Xiao Q, G. McPherson. 2009. *Testing a Bioswale to Treat and Reduce Parking Lot Runoff*. Center for Urban Forest Research. USDA Forest Service. Viewed online at: http://www.fs.fed.us/psw/programs/uesd/uep/products/psw_cufr761_P47ReportLRes_AC.pdf.
- Ziemer, RR. 1987. "Water yield from forests: an agnostic view: In: Callaham RZ, DeVries JJ, editors. *Proceedings of the California Watershed Management Conference*. California Watershed Management Conference. Nov. 18-20, 1986. West Sacramento (CA): University of California

Wildland Resources Center. pp. 74-78. Viewed online at:
<http://www.treesearch.fs.fed.us/pubs/8672>.

Ziemer RR, Lisle TE. 1998. "Hydrology." In: Naiman, RJ; Bilby, RE, editors. *River Ecology and Management: Lessons from the Pacific Coastal Ecoregion*. New York (NY): Springer-Verlag. Chapter 3. pp. 43-68. Viewed online at:
http://www.fs.fed.us/rm/boise/publications/misc/naiman_riparianecology.pdf.

Zinke PJ. 1987. "The recovery factor in watershed land use for timber harvest and growth." Impact '87. California Licensed Foresters Association (CLFA) Annual Conference. Mar. 6-7, 1987 Sacramento (CA): California Licensed Foresters Association. 7 pp.

Additional References

Battany MC, Grismer ME. 2000. "Rainfall runoff and erosion in Napa Valley vineyards: effects of slope cover and surface roughness." [Journal.] *Hydrological Processes*. 14:1289-1304.

Bayley PB, Li HW. 2008. "Stream fish response to grazing exclosures." [Journal.] *North American Journal of Fisheries Management*. 28:135-147.

Day SD, Dickinson SB, editors. 2008. *Managing Stormwater for Urban Sustainability using Trees and Structural Soils*. Blacksburg (VA): Virginia Polytechnic Institute and State University. Viewed online at: <http://www.vtfpr.org/urban/documents/TreesAndStructuralSoilsManual.pdf>. Accessed: Aug., 2011.

Diamond JM. 2005. *Collapse: How Societies Choose to Fail or Succeed*. New York (NY): Viking Press. 575 pp. Erman DC, Newbold JD, Roby KB. 1977. *Evaluation of Streamside Buffer Strips for Protecting Aquatic Organisms*. Davis (CA): Contribution No. 165. California Water Resources Center. University of California, Davis. 48 pp.

Giono J. 2005. *The Man Who Planted Trees*. 20th anniversary edition. White River Junction (VT): Chelsea Green. 61 pp.

Harris RR, Cafferata PH. 2005. Forest Loss and Fragmentation. Session 2: Impacts and Implications—Impacts on Water Quantity and Quality [Abstract]. PowerPoint presentation. California Forest Futures 2005 Conference. May 24- 25. Sacramento (CA):

Herbst DB, Kane JM. 2004. *Responses of Stream Channels Riparian Habitat and Aquatic Invertebrate Community Structure to Varied Livestock Grazing Exposure and Management in the West Walker River Watershed (Mono County California)*. South Lake Tahoe (CA): Lahontan Regional Water Quality Control Board. Final report submitted in partial fulfillment of Lahontan Regional Water Quality Control Board contract numbers 9-046-256-0 9-118-160-0 and 0-091-160-0. 41 pp. Viewed online at: http://www.swrcb.ca.gov/rwqcb6/water_issues/programs/swamp/docs/herbst_kane_wwgs_2004.pdf.

- Hymanson ZP, Collopy MW, editors. 2010. *An Integrated Science Plan for the Lake Tahoe Basin: Conceptual Framework and Research Strategies*. Albany (CA): USDA Forest Service. Pacific Southwest Research Station. Chapter 4.11. p. 117-123. General Technical Report PSW-GTR-226. Viewed online at: http://www.fs.fed.us/psw/publications/documents/psw_gtr226/psw_gtr226.pdf.
- Knapp RA, Matthews KR. 1996. "Livestock grazing, golden trout, and streams in the Golden Trout Wilderness, California: Impacts and management implications." [Journal.] *North American Journal of Fisheries Management*. 16:805-820.
- Madej MA, Ozaki V. 1996. "Channel response to sediment wave propagation and movement, Redwood Creek, California, USA." [Journal.] *USA Earth Surface Processes and Landforms*. 21(10):911-927.
- McGurk BJ, Berg NH, Davis ML. 1996. "Camp and Clear Creeks El Dorado County: predicted sediment production from forest management and residential development." In: *Sierra Nevada Ecosystem Project. Final Report to Congress, Volume II: Assessments and Scientific Basis for Management Options*. Davis (CA): University of California, Centers for Water and Wildland Resources. Chapter 53. pp. 1407-1420. Viewed online at: http://pubs.usgs.gov/dds/dds-43/VOL_II/VII_C53.PDF.
- McPherson EG, Simpson JR, Peper PJ, Xiao QF. 1999. *Tree Guidelines for San Joaquin Valley Communities*. Davis (CA): USDA Forest Service. Pacific Southwest Research Station. Western Center for Urban Forest Research and Education. 59 pp.
- McPherson E, Gregory S, James R, Peper P, Crowell J, Aaron MN, Xiao Q. 2010. *Northern California Coast Community Tree Guide: Benefits, Costs, and Strategic Planting*. Albany (CA): General Technical Report PSW-GTR-228. USDA Forest Service. Pacific Southwest Research Station.
- Miller WW, Johnson DW, Denton C, Verburg PSJ, Dana GL, Walker RF. 2005. "Inconspicuous nutrient laden surface runoff from mature forest Sierran watersheds." [Journal.] *Water, Air and Soil Pollution*. 163:3-17.
- Murphy DD, Stine PA, editors. 2004. *Proceedings of the Sierra Science Symposium*. General Technical Report PSW GTR-193. Oct. 7-10, 2002. Kings Beach (CA): USDA Forest Service. Pacific Southwest Research Station. Albany (CA): 287 pp.
- Newbold JD, Erman DC, Roby KB. 1980. "Effects of logging on macroinvertebrates in streams with and without buffer strips." [Journal.] *Canadian Journal of Fisheries and Aquatic Science*. 37:1076-1085.
- Ponce VM, Lindquist, DS. 1990. "Management of baseflow augmentation: a review." [Journal.] *Water Resources Bulletin*. 26(2):259-268.
- University of California, Centers for Water and Wildland Resources. 1996. *Sierra Nevada Ecosystem Project. Final Report to Congress. Volume I. Assessment Summaries and Management Strategies*.

Davis (CA): Chapter 8. Watersheds and Aquatic Biodiversity. pp. 123-133. Viewed online at:
http://pubs.usgs.gov/dds/dds-43/VOL_I/VI_C08.PDF.

USDA Forest Service. 2002. *Control Stormwater Runoff with Trees*. Davis (CA): Fact Sheet #4. USDA Forest Service. Pacific Southwest Research Station. Center for Urban Forest Research. 2 pp.